

Anti-protozoan activities of *Harungana madagascariensis* stem bark extract on trichomonads and malaria

E.O. Iwalewa, N.O. Omisore, C.O. Adewunmi, A.A. Gbolade, O.G. Ademowo, C. Nneji, O.I. Agboola, O.M. Daniyan

Department of Pharmacology, Faculty of Pharmacy, Obafemi Awolowo University, Ile-Ife, Nigeria

Department of Drug Research and Production Unit, Faculty of Pharmacy, Obafemi Awolowo University, Ile-Ife, Nigeria

Department of Pharmacognosy, Faculty of Pharmacy, Obafemi Awolowo University, Ile-Ife, Nigeria

Institute of Medical Research and Training, College of Medicine, University of Ibadan, Ibadan, Nigeria

Department of Paraclinical Sciences (Phytomedicine Programme), Faculty of Veterinary Sciences, University of Pretoria, PB 0X4, Onderstepoort 0110, Pretoria, South Africa

With 5 tables and 2 figures

Abstract

Aim of the study: The ethanolic stem bark extract of *Harungana madagascariensis* (Hypericaceae), (Choisy) Poir were evaluated for their activities on *Trichomonas gallinae* (Rivolta) Stabler isolated from the pigeon (*Columba livia*). It was also tested for their anti-malarial activity on N67 *Plasmodium yoelii nigeriensis* (*in vivo*) in mice and on *Plasmodium falciparum* isolates *in vitro*.

Materials and methods: The anti-trichomonal screening was performed *in vitro* using *Trichomonas gallinae* culture. The minimum lethal concentration (MLC) is the lowest concentration of the test extract in which no motile organisms were observed. The anti-malarial effects were determined *in-vivo* for suppressive, curative and prophylactic activities in mice receiving a standard inoculum size of 1×10^7 (0.2 ml) infected erythrocytes of *Plasmodium yoelii nigeriensis* intraperitoneally, and the *in vitro* was performed against 3 isolates of *Plasmodium falciparum* in a candle jar procedures.

Results: The IC₅₀ of the extract and metronidazole (MDZ) (Flagyl) on *Trichomonas gallinae* at 48 h are 187 and 1.56 µg/ml. The IC₅₀ of the extract, chloroquine (CQ) and artemether (ART) on *Plasmodium falciparum* are between 0.052 and 0.517 µg/ml for the extract and 0.021 and 0.0412 µg/ml for ART and CQ, respectively. The actions of the extract in *in vivo* study on *Plasmodium yoelii nigeriensis* showed that in both suppressive and prophylactic tests the percentages chemo-suppressive were between 28.6–44.8% and 30.2–78.2% respectively, while only 80 mg/kg of the extract reduced the parasitaemia level when compared to the control and the standard drugs in curative test.

Conclusions: *Harungana madagascariensis* stem bark extract therefore exhibited significant anti-protozoan effects against *Trichomonas* and *Plasmodium* both *in vivo* and *in vitro*.

Keywords: Anti-malarial; Anti-trichomonal; *Harungana madagascariensis*; Stem bark extract; *In vivo* and *in vitro* studies

1. Introduction

1.1. Plant

Harungana madagascariensis (Choisy) Poir. (Hypericaceae) is otherwise called dragon's blood tree. The genus is monotypic, the single species being found in tropical Africa, Madagascar, Mauritius and Mascareigne Island. The local names are "Amuje", "Aroje", and "Arunje". It is widespread and locally abundant in areas where annual rainfall exceeds 1300mm. (Hutchinson and Dalziel, 1954; Humphries and Stanton, 1992; Csurhes and Edwards, 1998). The fresh stem barks of *Harungana madagascariensis* were collected along

Ede and Ondo roads near the main University campus, Ile-Ife. The plant was identified by Mr. O.A. Oladele of the Department of Pharmacognosy, Faculty of Pharmacy, Obafemi Awolowo University, Ile-Ife and a voucher specimen with voucher number FHI 107392 was kept at the herbarium of the Forestry Research Institute of Nigeria, Ibadan.

1.2. Previously isolated classes of constituents

Some of the constituents and isolated compounds from *Harungana madagascariensis* includes: flavonoids, alkaloids, saponins, glycosides, and tannins (Moulari et al., 2006a,b). Three prenylated anthranoids: harunmadagascariens C and D and kenganthranol D, three prenylated 1,4-anthraquinone: harunmadagascariens A and B, harunganol B and harungin anthrone (Kouam et al., 2006a,b, 2007), Bazouanthrone, a newanthrone derivative, has been isolated from the root bark of *Harungana madagascariensis*, together with other known compounds, feruginin A, harunganin, harunganol A, harunganol B, friedelan-3-one and betulinic acid (Ndjakou Lenta et al., 2007).

1.3. Uses in traditional medicine

Traditionally, the leaves and stem bark are used for the treatment of anaemia, the stem bark is also used for nephrosis, malaria, gastro-intestinal disorders and fever (EMEA, 1999; Tona et al., 2000; Erah et al., 2003; Kamanzi Atindehou et al., 2004). Aqueous leaf extract has been shown to have anti-microbial activity on different strains of bacteria (*Bacillus subtilis*, *Staphylococcus aureus*, *Eschericia coli* and *Salmonella typhi*), thus substantiating its use for gastro-intestinal disorders (Okoli et al., 2002). The stem bark extract inhibited *Entamoeba histolytica* growth at a concentration of less than 10µg/ml (Tona et al., 2000). The stem bark extract also exhibited more than 70% inhibition of acetylcholine and potassium chloride solution-induced contractions on isolated guinea-pig ileum. *Harungana madagascariensis* is a component of Jubi Formula, an herbal preparation which was found to restore the pack cell volume (PCV) and haemoglobins (Hb) concentration in anaemia conditions and is a potential substitute for blood transfusion (Erah et al., 2003). Lukwa et al. (2001) made an ethnobotanical survey on some herbal remedies used by traditional healers in Mola, Kariba district of Zimbabwe for the treatment and prevention of malaria, *Harungana madagascariensis* is one of the plants used for the treatment of malaria. Earlier studies of *Harungana madagascariensis* were based on the leaves, (Madubunyi et al., 1995; Kamanzi Atindehou et al., 2004; Okoli et al., 2002; Moulari et al., 2006a,b; Kouam et al., 2007). Nwodo (1989) and Kouam et al. (2006a,b) have, however, examined the effect of crude extracts and isolated compounds from a hexane extract of the stem bark of *Harungana madagascariensis*, for their analgesic, anti-inflammatory, alpha-glucosidase inhibition and anti-oxidant activities respectively. The most recent work of Ndjakou Lenta et al. (2007) showed the anti-plasmodial effects of six isolated compounds from the root bark of *Harungana madagascariensis in vitro*. Many isolated compounds from natural products have been tested *in vitro* for anti-malarial properties; and found to exhibit potent activities (Rukunga and Simons, 2006). Studies have shown that *in vivo* examinations of the same compounds are without such activities (Khalid et al., 1989; Bray et al., 1990). To complement these studies, we therefore, investigated the activities of *Harungana madagascariensis* stem bark on *Trichomonas gallinae (in vitro)* and malaria (*in vivo* and *in vitro*), in order to authenticate its efficacy in traditional medicine.

2. Materials and methods

2.1. Plant materials and preparation of the extracts

The ethanolic extraction was made by soaking 325 g of the dried powdered stem bark in 70:30 ethanol:water for 24 h. After maceration, the extract was evaporated *in vacuo* in a rotary evaporator to dryness, yielding 9 g of the brown crude extract.

2.2. Parasites

Trichomonas gallinae was obtained from the gut of a local pigeon bred and maintained in the Drug Research and Production Unit. The N 67 chloroquine (CQ) sensitive strain of *Plasmodium yoelii nigeriensis* was donated generously by the Malaria Research and Reference Reagent Resource Center (MR4) Manassas, VA, USA, and was obtained from the Institute of Medical Research and Training courtesy of Dr. O.G. Ademowo. The *Plasmodium*

falciparum was obtained from the patients that visited Clinical Pharmacology Unit, University Teaching Hospital, College of Medicine, Ibadan.

2.3. Animals

Mice of either sex, weighing between 18 and 220 g were used. The animals were maintained at 25 + 1 °C under natural 12 h daylight/night conditions for at least 5 days before the experiment. All the animals were fed with standard diet in the Department of Pharmacology Animal House and water was given *ad libitum*.

The "principle of laboratory animal care" (National Institute of Health-NIH publication No. 85-23) guidelines and procedures were followed in this study (NIH publication revised, 1985).

2.4. Anti-trichomonal test

Trichomonas gallinae was cultured in Ringer-Egg-Serummedium according to the method of Boeck and Drbolav (1925) as cited by Levine (1961). The anti-trichomonal screening was performed *in vitro* according to the method of Meingasser and Thurner (1979) as reported by Oyedapo et al. (2004). *Trichomonas* culture was distributed into micro well plates and solutions of the test substances were added to make concentrations of 250, 500, 1000, 2000 and 4000 µg/ml. Metronidazole (MDZ) was used as reference substance in the following concentrations: 0.62, 1.25, 2.5, 5, 10 and 20 µg/ml. 50 µl of extract and 150 µl of the culture solution were pipetted into the micro wells and incubated at 37 °C for 24 and 48 h. Wet mount microscopic observation was done for each time set. The experiments were done in triplicates. The minimum lethal concentration (MLC) is the lowest concentration of the test extract in which no motile organisms were observed. A graph of % inhibition of parasites growth against the concentration of the extract and the standard agents were plotted and the IC₅₀ (the inhibitory concentration that suppressed the growth of the parasites population by 50%) were extrapolated.

2.5. Anti-malarial tests

The anti-malarial effects were determined according to Peters and Robinson (1992), Ryley and Peters (1970) and Peters (1965), respectively for suppressive, curative and prophylactic tests, and the *in vitro* was performed according to WHO (1987).

The evaluation of blood schizonticidal activity on an early suppressive infection (4-day test), prophylactic and curative tests. The blood schizonticidal activity of the extracts (20, 40 and 80 mg/kg/day); chloroquine (5 mg/kg/day) as standard drug and normal saline (0.9%) as control were assessed in parasitized albino mice receiving a standard inoculum size of 1×10^7 (0.2 ml) infected erythrocytes of *Plasmodium yoelii nigeriensis* intraperitoneally on day zero. In the suppressive test, the administration of drugs took place from Day 0 to Day 3. On Day 4, blood smears were made from the tail of the mouse and percentage parasitemia determined. In prophylactic test, another group of animals were first treated with the extracts and pyrimethamine (1.2 mg/kg) daily for 3 days, and were inoculated with parasites after treatment, 72 h after infection, parasitaemia were determined. However in curative test, the animals were infected with same parasites and were allowed to be established for 72 h after which treatment was given for 5 days and parasitaemia was determined.

2.6. In vitro method

The extract (10mg) was weighed and dissolved in 0.5 ml of sterile distilled water to give a stock concentration of 5mg/ml. Chloroquine and artemether (ART) stock concentration were 0.5 mg/ml. From these stock solutions, a 10-fold dilution was made with RPMI 1640 medium (without bicarbonate). 100 µl aliquots of diluted extract dispensed into 96-well plates gave 0.07, 0.21, 0.62, 1.85, 5.60, 16.7, and 50.0 µg/ml as final concentration, while CQ and ART gave 10 times lower than that of the extract. Fifty microlitres of RPMI 1640 medium plus bicarbonate and serum A₊ as complete serum were aliquoted into all the wells. Twenty-five microlitres of infected red cell of each isolates were separately added. The starting parasitaemia was between 2.5–5.0%. However, the negative controls were without the extract. After this, the plates were covered, shaken gently and incubated in a desiccator at 37 °C for 27–30 h. After incubation, the contents of both the controls and tested wells were harvested and the deposited red cells were transferred to a slide to form a thick film. The film

was dried and stained with Giemsa and the parasites were assessed for growth. The numbers of schizonts with three or more nuclei out of a total of 200 asexual parasites were noted. A graph of %inhibition of parasites growth against the concentration of the extract and the standard agents were also plotted and the IC₅₀ were determined.

2.7. Data and statistical analysis

Data are expressed as mean ± S.E.M. and analyzed using one way analysis of variance (ANOVA) followed by Dunnett Test for comparing pairs of data. The significant level was set at $p < 0.05$. The IC₅₀ values were calculated using the Microsoft Excel program.

3. Results

Results of the anti-protozoan activity of HMSE on trichomonads and malaria parasites showed an inhibitory effect on the parasites growth. The *in vitro* activities were expressed in Tables 1 and 2; Figs. 1 and 2. The IC₅₀ of the extract and MDZ (Flagyl) on *Trichomonas gallinae* at 24 and 48 h are 1000.0 and 1.87 µg/ml at 24 h and 187.0 and 1.56 µg/ml at 48 h, respectively. The IC₅₀ of the extract, CQ and ART on *Plasmodium falciparum* are between 0.052 and 0.517 µg/ml for the extract and 0.021 and 0.0412 µg/ml for ART and CQ respectively. The actions of the extract in *in vivo* study on *Plasmodium yoelii nigeriensis* were indicated in Tables 3–5. In both suppressive and prophylactic tests the percentage chemo-suppressive was between 28.6–44.8% and 30.2–78.2%, respectively. However, in curative test, only 80mg/kg of the extract reduced the parasitaemia level as compared to the control and the standard drugs.

4. Discussion

Plants are often considered to be useful against infectious diseases, which are used as alternatives to synthetic compounds without regard to different etiological agents (viral, bacterial, fungus and protozoa) in traditional medical practice. The wide reports of *Harungana madagascariensis* plant parts that are used in various infectious disorders points to the fact that the plant constituents could be potential anti-infective candidates for the treatment of viral, bacteria, fungi, and protozoa (Madubunyi et al., 1995; Okoli et al., 2002; Kamanzi Atindehou et al., 2004; Moulari et al., 2006a,b; Kouam et al., 2007).

Our experiments demonstrate that phyto-chemical constituents present in the stem bark extract of *Harungana madagascariensis* used in traditional medicine to treat infectious diseases, exhibited anti-protozoan activity against *Trichomonas gallinae in vitro* and *Plasmodium* parasites both *in vivo* and *in vitro*. This however, supports the reports of Gessler et al. (1995), Tona et al. (2000) and Ndjakou Lenta et al. (2007) that showed the leaves and root bark extracts from *Harungana madagascariensis*, which inhibited another protozoan *Entamoeba histolytica* growth in an investigation on the putative anti-amoebic and *Plasmodium falciparum* isolates respectively. Although, anti-protozoan effects of different parts (leaves, seeds and root extracts) of *Harungana madagascariensis* have been evaluated by various authors (Tona et al., 2000; Kamanzi Atindehou et al., 2004 and Ndjakou Lenta et al., 2007), no activity on the stem bark or any part of the plant have been reported *in vivo*. All the works cited are *in vitro* studies. In addition, other forms of protozoa (amoebiasis, leishmaniasis, trypanosomiasis) have been tested against different extracts of this plant, while trichomonads remain the only protozoa. These two reasons prompted the objective of this study.

Trichomoniasis and malaria have important medical, social and economic implications. Premature labour, premature ruptures of placental membranes and low birth-weights have been reported in infected pregnant women. *T. vaginalis* can augment the predisposition of individuals to human immunodeficiency viruses (HIV) (Petrin et al., 1998). Metronidazole is the recommended treatment for trichomoniasis. Tinidazole (Fasigyn) is an alternative drug. Adverse reactions to high doses and also allergic reactions to metronidazole have been reported (Narcisi and Secor, 1996). Multiple drug resistances to these routine drugs are on the increase, it has been shown that 5% of the population has shown resistance to metronidazole (Lossick, 1989) and the frequency of chloroquine resistance ranged from 18% to 62% (Rowland et al., 1997).

This problem of resistance alone is enough to search for new agents from plants used in traditional medicine for the treatment of these ailments (Rukunga and Simons, 2006). The activity of the extract on trichomonads is weak compared to metronidazole. However, the IC₅₀ of the extract at 48 h was highly reduced; an indication that the extract contact time with the parasite is a factor for its effectiveness. On malaria parasites also, the IC₅₀ of the extract was highly comparable with that of chloroquine and artemether, however, the positive control drugs are 10 times more potent than the extract against the two parasites tested. Rukunga and Simons (2006) classified the *in vitro* activity of natural products having anti-plasmodial effects into three categories. The first category deals with plant extracts exhibiting high anti-plasmodial action (IC₅₀ less than or equal to 10 µg/ml). The second groups are moderately active with IC₅₀ greater than 10 but less than 100 µg/ml, while the third category was termed as low active group as their IC₅₀ is greater than 100 µg/ml. The results of this study fall in the first category indicating that the extract has a high anti-plasmodial activity against malaria, but exhibited weak effect on trichomonads. These observations indicated that a difference in the parasites genus has an effect on the response to the test substance. In the three models of malaria treatment employed in our *in vivo* study, the results showed that the extract also inhibited the multiplication of malaria parasites in rodents in various degrees. It was clearly shown that the extract had no significant effect in curative treatment, but is highly effective through suppressive and prophylactic mode of therapy. This could therefore be the mode of treatment employed in traditional medicine. Many compounds from natural products have been shown to exhibit potent *in vitro* anti-malarial activities (Rukunga and Simons, 2006). However, various studies have indicated that *in vivo* examinations of the same compounds are without such activities (Khalid et al., 1989 and Bray et al., 1990). Till date no single report is available on the *in vivo* effect of the stem bark extract against *Plasmodium* in rodents. The current study reports for the first time, the *in vivo* efficacy of the plant to justify the traditional uses against malaria and *Trichomonas gallinae* as anti-protozoal agent. These activities on malaria and trichomonads pathogens like those reported against fungi and bacteria organisms could then explain the usefulness of the plant in the treatment of fever, tuberculosis, diarrhea, dysentery, syphilis, gonorrhoea, parasitic skin diseases and wounds. From the phytochemical studies, anthraquinone, anthranoids, coumarins, and triterpenoids are the compounds that were detected. The Hypericaceae family is well known for the production of various phenolic compounds such as anthraquinones, xanthenes, coumarins, biflavonoids and anthrone derivatives (Iinuma et al., 1995) which are known to exhibit antimicrobial activities (Rabanal et al., 2002). It is possible that these bioactive compound(s) found in the stem bark could be responsible for these activities.

5. Conclusion

Harungana madagascariensis stem bark extract had significant anti-trichomonal and anti-malarial activities thus indicating their positive role, and justifying the ethnomedicinal use of the plant parts in traditional medicine.

References

1. Boeck, W.C., Drbolav, J., 1925. Laboratory culture of protozoal parasites. Proceedings of National Academy of Sciences 11, 235–238.
2. Bray, D.H., Warhurst, D.C., Connolly, J.D., O'Neill, M.J., Phillipson, J.D., 1990. Plants as sources of antimalarial drugs. Part 7. Activity of some species of meliaceae plants and their constituent limonoids. Phytotherapy Research 4, 29–35.
3. Csurhes, S., Edwards, R., 1998. Potential Environmental Weeds in Australia: Candidate Species for Preventative Control. Queensland Department of Natural Resources, p. 109.
4. EMEA, 1999. Committee for Veterinary Medicinal Products – *Harungana madagascariensis* Summary Report – The European Agency for the Evaluation of Medicinal Products, Veterinary Medicines Evaluation Units. EMEA/MRL/598/99-FINAL.
5. Erah, P.O., Asonye, C.C., Okhamafe, A.O., 2003. Response of *Trypanosoma brucei brucei*-induced anaemia to a commercial herbal preparation. African Journal of Biotechnology 2, 307–311.

6. Gessler, M.C., Tanner, M., Chollet, J., Nkunya, M.H.H., Heinrich, M., 1995. Tanzanian medicinal plants used traditionally for the treatment of malaria: in vivo antimalarial and in vitro cytotoxic activities. *Phytotherapy Research* 9, 504–508.
7. Humphries, S.E., Stanton, J.P., 1992. Weed assessment in the wet tropics world heritage area of north Queensland. Report to the Wet Tropics Management Agency, pp. 25–26.
8. Hutchinson, J., Dalziel, J.M., 1954. *Flora of West Tropical Africa*, Part 1, vol. 1. Crown Agents, London, p. 290.
9. Iinuma, M., Hideki, T., Tetsuro, I., Toshiyuki, T., Mohammad, A., 1995. Two prenylated anthrones in *Harungana madagascariensis*. *Phytochemistry* 40, 267–270.
10. Levine, N.D., 1961. Laboratory diagnosis of protozoan infections. In: *Protozoan Parasites of Domestic Animals and Man*. Burgess Publishing Co., Minneapolis, pp. 377–393.
11. Lossick, J.G., 1989. Therapy of urinogenital Trichomoniasis. In: Honigberg, B.M. (Ed.), *Trichomonads Parasitic in Man*. Springer Verlag, New York, pp. 324–341.
12. Lukwa, N., Mutambu, S.L., Makaza, N., Molgaard, P., Furu, P., 2001. Perceptions about malaria transmission and control using anti-malaria plants in Mola, Kariba, Zimbabwe. *Nigerian Journal of Natural Products and Medicine* 5, 4–7.
13. Kamanzi Atindehou, K., Schmid, C., Brun, R., Kone, M.W., Traore, D., 2004. Antitrypanosomal and antiplasmodial activity of medicinal plants from Côte d'Ivoire. *Journal of Ethnopharmacology* 90, 221–227.
14. Khalid, S.A., Duddeck, H., Gonnzalez-Sierra, M., 1989. Isolation and characterization of an antimalarial agent of the neem tree *Azadirachta indica*. *Journal of Natural Products* 52, 922–926.
15. Kouam, S.F., Khan, S.N., Krohn, K., Ngadjui, B.T., Lapche, D.G., Yapna, D.B., Zareem, S., Moustafa, A.M., Choudhary, M.I., 2006a. Alpha-glucosidase inhibitory anthranols, kenganthranols A–C, from the stem bark of *Harungana madagascariensis*. *Journal of Natural Product* 69, 229–233.
16. Kouam, S.F., Ngadjui, B.T., Krohn, K., Wafo, P., Ajaz, A., Choudhary, M.I., 2006b. Prenylated anthronoid antioxidants from the stem bark of *Harungana madagascariensis*. *Brazilian Journal of Medical and Biological Research* 38, 1087–1094.
17. Kouam, S.F., Yapna, D.B., Krohn, K., Ngadjui, B.T., Ngoupayo, J., Choudhary, M.I., Schulz, B., 2007. Antimicrobial prenylated anthracene derivatives from the leaves of *Harungana madagascariensis*. *Journal of Natural Products* 70, 600–603.
18. Madubunyi, I.I., Obi, S.K.C., Nwebube, N.I., Chime, A.B., 1995. Antihepatotoxic and antimicrobial activities of *Harungana madagascariensis* leaf extracts. *International Journal of Pharmacognosy* 33, 129–134.
19. Meingasser, J.G., Thurner, J., 1979. Strain of *Trichomonas vaginalis* resistant to metronidazole and other 5-nitroimidazoles. *Antimicrobial Agents and Chemotherapy* 15, 254–257.
20. Moulari, B., Lboutounne, H., Chaumont, J.-P., Guillaume, Y., Millet, J., Pellequer, Y., 2006a. Potentiation of the bactericidal activity of *Harungana madagascariensis* Lam. ex Poir. (Hypericaceae) leaf extract against oral bacteria using poly(D,L-lactide-co-glycolide) nanoparticles: in vitro study. *Acta Odontologica Scandinavica* 64, 153–158.
21. Moulari, B., Pellequer, Y., Lboutounne, H., Girard, C., Chaumont, J.-P., Millet, J., Muyard, F., 2006b. Isolation and in vitro antibacterial activity of astilbin, the bioactive flavanone from the leaves of *Harungana madagascariensis* Lam. ex Poir. (Hypericaceae) *Journal of Ethnopharmacology* 106, 272–278.
22. Narcisi, E.M., Secor, W.E., 1996. In vitro effect of Tinidazole and Furazolidone on metronidazole-resistant *Trichomonas vaginalis*. *Antimicrobial Agents Chemotherapy* 40, 1121–1125.
23. Ndjakou Lenta, B., Ngouela, S., Fekam Boyom, F., Tantangmo, F., Feuya Tchouya, G.R., Tsamo, E., Gut, J., Rosenthal, P.J., Donald Connolly, J., 2007. Anti-plasmodial activity of some constituents of the root bark of *Harungana madagascariensis* LAM. (Hypericaceae) *Chemical and Pharmaceutical Bulletin* 55, 464–467.
24. Nwodo, O.F.C., 1989. Antibiotic and anti-inflammatory analgesic activities of *Harungana madagascariensis* stem bark. *International Journal of Crude Drug Research* 27, 137–140.

25. Okoli, A.S., Okeke, M.I., Iroegbu, C.U., Ebo, P.U., 2002. Antibacterial activity of *Harungana madagascariensis* leaf extract. *Phytotherapy Research* 16, 174–179.
26. Oyedapo, O.A., Adewunmi, C.O., Makanju, V.O., Iwalewa, E.O., Adenowo, T.K., 2004. Antitrichomonal activities of synthetic 1,3-diaryl-2-propen-1-ones on *Trichomonas gallinae*. *African Journal of Traditional, Complimentary and Alternative Medicine* 1, 55–62.
27. Peters, W., 1965. Drug resistance in *Plasmodium berghei*. Venke and Lips (1948). I. Chloroquine resistance. *Experimental Parasitology* 17, 80–87.
28. Peters, W., Robinson, B.L., 1992. The chemotherapy of rodent malaria. XLVII. Studies on pyronaridine and other Mannich base antimalarials. *Annal Journal of Tropical Medicine and Parasitology* 86, 455–465.
29. Petrin, D., Delgaty, K., Bhatt, R., Garber, G., 1998. *Clinical Microbiology Reviews* 11, 300–317.
30. Rabanal, R.M., Arias, A., Prado, B., Herna'ndez-Pe' rez, M., Sa' nchez-Mateo, C.C., 2002. Antimicrobial studies on three species of *Hypericum* from the Canary Islands. *Journal of Ethnopharmacology* 81, 287–292.
31. Rowland, M., Durrani, N., Hewitt, S., Sondorp, E., 1997. Resistance of falciparum malaria to chloroquine and sulfadoxine-pyrimethamine in Afghan refugee settlements in western Pakistan: surveys by the general health services using a simplified in vivo test. *Tropical Medicine of International Health* 2, 1049–1056.
32. Rukungu, G., Simons, A.J., 2006. The Potential of Plants as a Source of Antimalarial Agents-A Review. *Africa Herbal Antimalarial Meeting CDE and ICRAF, Nairobi*, p. 9.
33. Ryley, J.F., Peters, W., 1970. The antimalarials of some quinoline esters. *Annal Journal of Tropical Medicine and Parasitology* 84, 209.
34. Tona, L., Kambu, K., Ngimbi, N., Mesia, K., Penge, O., Lusakibanza, M., Cimanga, K., De Bruyne, T., Apers, S., Totte, J., Pieters, L., Vlietinck, A.J., 2000. Antiamoebic and spasmolytic activities of extracts from some antidiarrhoeal traditional preparations used in Kinshasa, Congo. *Phytomedicine* 7, 31–38.
35. World Health Organization, 1987. In vitro micro test (mark II) for the assessment of the response of *Plasmodium falciparum* to Chloroquine Mefloquine, Quinine, Sulphadoxine-Pyrimethamine and Amodiaquine. W.H.O. mimeographed doc. No. MAP/87.2.

Table 1

An *in vitro* anti-trichomonads activity of metronidazole (MDZ) as positive control, and the stem bark (HMSE) extract of *Harungana madagascariensis* on *Trichomonas gallinae* (24 and 48 h)

Percentage of the number of surviving trichomonads (%mortality/inhibition of <i>Trichomonas gallinae</i> growth)				
Concentration ($\mu\text{g/ml}$)	24 h		48 h	
	Flagyl MDZ	HMSE	Flagyl MDZ	HMSE
0	0.0	0.0	0.0	0.0
0.62	11.1 \pm 5.5	–	28.2 \pm 1.3	–
1.25	25.1 \pm 4.6	–	38.0 \pm 1.9	–
2.5	94.4 \pm 2.5	–	95.5 \pm 3.6	–
5.0	100.0	–	100.0	–
10	100.0	–	100.0	–
20	100.0	–	100.0	–
250	ND	14.6 \pm 2.2	ND	66.2 \pm 1.5
500	ND	33.4 \pm 2.6	ND	76.1 \pm 0.7
1000	ND	48.1 \pm 3.8	ND	84.3 \pm 0.9
2000	ND	88.8 \pm 0.9	ND	97.6 \pm 0.3
4000	ND	100.0	ND	100.0

ND: not determined.

Table 2

The anti-trichomonads and anti-malarial IC50 values of the extract of *Harungana madagascariensis* and standard drugs-metronidazole, chloroquine and artemether on both trichomonads and malarial parasites (*in vitro*)

IC ₅₀ of HMSE and MDZ on <i>Trichomonads gallinae</i> at 24 and 48 h (μ g/ml)			
24 h		48 h	
HMSE	MDZ	HMSE	MDZ
1000.00	1.87	187.00	1.56

IC ₅₀ of HMSE, CQ and ART on <i>Plasmodium falciparum</i> (μ g/ml)	
HMSE	0.052–0.517
CQ	0.0412
ART	0.021




Fig. 1. *In vitro* effects of *Harungana madagascariensis* extract on *Plasmodium falciparum*. %Inhibition of parasite growth against the various concentrations of the stem bark extract of *Harungana madagascariensis* (HMSE) on *Plasmodium falciparum* isolates (◆-◆) 001, (■-■) 011, (▲-▲) 017




Fig. 2. *In vitro* effects of chloroquine (CQ) and artemether (ART) against *Plasmodium falciparum* isolates. %Inhibition of parasite growth against the various concentrations of the standard drugs chloroquine (CQ) (■-■) and artemether (ART) (▲-▲) on *Plasmodium falciparum* isolate.

Table 3

The suppressive (*in vivo*) activity of the stem bark (HMSE) extract of *Harungana madagascariensis* and chloroquine (CQ) on *Plasmodium yoelii nigeriensis*

Suppressive test		
	%Parasitaemia	%Chemo-suppression
Control	34.65 ± 1.31	–
20 mg/kg	24.75 ± 2.25 ^{a,b}	28.57
40 mg/kg	21.84 ± 1.61 ^{a,b}	36.99
80 mg/kg	19.14 ± 1.07 ^{a,b}	44.76
CQ (5 mg/kg)	10.2 ± 0.3 ^a	70.56

Values are mean ± S.E.M., *n*=5; **p* < 0.05 vs. control.

^a Significantly different from control at *p* < 0.05.

^b Significantly different from CQ at *p* < 0.05.

Table 4

The prophylactic (*in vivo*) activity of the stem bark (HMSE) extract of *Harungana madagascariensis* and pyrimethamine (PYR) on *Plasmodium yoelii nigeriensis*

Prophylactic test		
	%Parasitaemia	%Chemo-suppression
Control	15.35 ± 0.12	–
20 mg/kg	10.71 ± 0.81 ^{a,b}	30.22
40 mg/kg	4.96 ± 0.83 ^{a,b}	67.69
80 mg/kg	3.35 ± 0.77 ^{a,b}	78.16
Pyrimethamine (1.2 mg/kg)	8.70 ± 0.53 ^a	43.32

Values are mean ± S.E.M., *n*=5; **p* < 0.05 vs. control.

^a Significantly different from control at *p* < 0.05.

^b Significantly different from PYR at *p* < 0.05.

Table 5

The curative (*in vivo*) activity of the stem bark (HMSE) extract of *Harungana madagascariensis* and chloroquine (CQ) on *Plasmodium yoelii nigeriensis*

	Curative test (%Parasitaemia)		
	D7	D9	D11
Control	13.5 ± 0.88	21.65 ± 1.41	22.12 ± 2.75
20 mg/kg	17.10 ± 1.09 ^{a,b}	21.78 ± 1.52	32.0 ± 1.36 ^{a,b}
40 mg/kg	15.94 ± 1.41 ^b	19.71 ± 1.24 ^a	25.92 ± 1.57 ^b
80 mg/kg	2.02 ± 0.43 ^{a,b}	12.63 ± 1.64 ^{a,b}	10.15 ± 0.85 ^{a,b}
CQ (5 mg/kg)	9.17 ± 1.21 ^a	18.10 ± 0.52	16.59 ± 1.10 ^a

Values are mean ± S.E.M., *n*=5; **p* < 0.05 vs. control.

^a Significantly different from control at *p* < 0.05.

^b Significantly different from CQ at *p* < 0.05.