

# PRELIMINARY PHYTOCHEMICAL SCREENING AND ANTIPROLIFERATIVE EFFECTS OF METHANOLIC EXTRACT OF STEM BARK OF DIOSPYROS MESPILIFORMIS HOCHST (EBENACEAE)

<sup>1</sup>ANAS ABBA, <sup>2</sup>ABDULKARIM AGUNU, <sup>3</sup>ABUBAKAR AHMED, <sup>4</sup>YAHYA IBRAHIM, <sup>5</sup>USMAN MOHAMMAD JAJERE, <sup>6</sup>UMAR SULEIMAN ABUBAKAR, <sup>7</sup>ABUBAKAR MOHAMMED KABIR

<sup>1,2,3,7</sup>Department of Science Laboratory Technology, College of Science and Technology, Hussaini Adamu Federal Polytechnic, P.M.B; 5004, Kazaure, Jigawa State, Postal code;234, Nigeria.

<sup>1,2,3</sup>Department of Pharmacognocny and Drug Development, Faculty of pharmaceutical sciences, Ahmadu Bello University Zaria, Kaduna, Nigeria.

<sup>4,5,6</sup>Department of Pharmacognocny and Drug Development, Faculty of pharmaceutical sciences, Ahmadu Bello University Zaria, Kaduna, Nigeria.

E-mail: <sup>1</sup>anasringim@yahoo.co.uk, <sup>2</sup>agunua@yahoo.com, <sup>3</sup>ahmedabubakar@abu.edu,

**Abstract**-According to ethnomedicinal information plant *Diospyros mespiliformis* Hochst (Ebenaceae) is commonly known as *Kanya* in Northern Nigeria which has been used widely in treating various ailments such as fever, whooping cough, wounds. Malaria, Pneumonia, Syphilis, Leprosy, Dermatomyces, Diarrhea without scientific validation. Preliminary Phytochemical screening, Thin layer chromatographic profile of methanol crude extract and antiproliferative studies were carried out in this research. Preliminary Phytochemical screening revealed the presence of Carbohydrate, Glycoside, Anthraquinone, Steroid, Triterpenes, Saponin, Tannins, Flavonoids and Alkaloid. TLC profile of the crude extract gave four spots with good  $R_f$  values. Antiproliferative evaluation were carried out using Guinea corn (*Sorghum bicolor*) seeds spread in a 9 cm wide petridish laid with cotton wool and Whatman filter paper which was treated with 1-30 mg/ml of methanolic extract in 24-96 hr period of incubation. At 24 hrs of incubation, the methanol extracts had  $24.771 \pm 0.526$  mm length of growth for the controls whereas the seeds treated with 10, 20, and 30 mg/ml of the extract produced a length total of  $2.772 \pm 0.494$  mm,  $2.150 \pm 0.490$  mm and  $2.257 \pm 0.489$  mm respectively, while at the end of 96 hours of incubation period, the radicles length of the control seeds measured  $93.77 \pm 9.730$  mm while those treated with 10, 20, and 30 mg/ml were observed to be  $37 \pm 3.297$  mm,  $17.023 \pm 2.802$  mm and  $16.086 \pm 1.976$  mm. This reduction in the growth implied 60.54, 81.87 and 82.83% respectively compared to the controls. All data were expressed as mean $\pm$ SEM and one way Analysis of Variance Anova statistical test using SAS Version 9.2 to test the significance.  $P < 0.05$  was considered Significance. This study has scientifically justified the traditional uses of *Diospyros mespiliformis* stem bark extracts for it has an antiproliferative property against radicles of a Guinea corn (*Sorghum bicolor*) which may relate to its use as anticancer agent. However, use of cancer cell lines will further confirm this claim.

**Keyword**-Antiproliferative effects , *Diospyros Mespiliformis*, Phytochemical,

## I. INTRODUCTION

*Diospyros mespiliformis* Hochst (Ebenaceae) has a fantastic mutualism and symbiotic network with many living organisms, from human beings to small insects. There is a complex ecological system revolving around this tree. It is one of the savanna giants that can live for more than 200 years. It is a tall, upright tree that can reach a height of 25 m, with a trunk circumference of more than 5 m. It has a dense evergreen canopy [1] The bark is black to grey, with a rough texture. The fresh inner skin of the bark is reddish. Leaves are simple, alternate, leathery and dark green. The margin is smooth and new leaves in spring are red, especially in young plants. Flowers are cream-coloured and bell-shaped [1]. Among Hausa Fulani people of Jigawa in Nigeria, *Diospyros mespiliformis* Hochst (Ebenaceae) happens to be one of such plant used in the treatment of tumor-related disease. In local Nigerian language (*Hausa*) the plant is known as *Kanya*. This very plant is not used only to treat tumor related disease but also to treat diseases such as fever, malaria, pneumonia, syphilis, leprosy,

dermatomyces [2]. It is also used in the treatment of diarrhea, whooping cough, wounds [3].

## II. MATERIAL AND METHOD

### *Collection and authentication of plant material*

The plant specimen for the studies were collected from Ringim local government of Jigawa State, Nigeria in January 2013. It was authenticated by a Taxonomist Muhammad Umar Galla of Biological Sciences Department, Ahmadu Bello University Zaria with voucher specimen number 901431.

### *Extraction of plant materials*

About 200g of powdered plant material were macerated in a separating funnel with 600ml of Methanol for one day (24hr) at room temperature with occasional shaking. The content was then filtered with cotton plug. The filtrate were then concentrated to dryness using a water bath at 60°C.

### **Preliminary Phytochemical Studies**

This procedure was carried out on the Methanol extracts according to [4],[5],[6] and [7] as outlined below.

**Molisch test;** to 2 ml of the extract in a test tube, few drops of molisch reagent and sulphuric acid was added and the colour reactions was recorded.

**Fehling test;** 5 ml of fehling solution A and B was added to 2 ml of the extract in a test tube and the colour reaction was recorded. This was passed over hot water bath for 15 minutes and the result was also observed and recorded.

**Ferric chloride test;** 0.5 ml of the extracts were dissolved in 10 ml of water each and filtered. Few drops of ferric chloride were added to the filtrate and the colour reaction was observed and recorded.

**Lead sub-acetate test;** 3 drops of lead sub-acetate solution was added to the extract solution and reaction was observed and recorded.

**Frothing test;** about 2 ml of the extract was dissolved in 10 ml of water and shaken vigorously for 30 seconds and allowed to stand for 30 minutes before observing and recording the reaction.

**Lieberman-Burchard test;** 1 ml of acetic anhydride was added to 1 ml of the extract. Few drops of sulphuric acid were carefully then added to the solution above and the reaction was observed and recorded.

**Salkowski test;** 2 ml of chloroform and few drops of sulphuric acid were added to about 2 ml of the extract and the reaction was observed and recorded.

**Bontrager test;** 2 ml of the extracts was added to 10 ml of benzene and shaken. This was then filtered and 5 ml of 10% ammonia solution was added to the filtrate and stirred and the reaction was observed and recorded.

**Shinoda test;** about 0.5 g of the extract was dissolved in 2 ml of 50% methanol. Few drops of magnesium filings and 3 drops of hydrochloric acid were added and the reaction observed and recorded.

**Sodium hydroxide test;** few drops of sodium hydroxide was added to 5 ml of the extract and the reaction was observed and recorded.

**Keller-killiani test;** 2 ml of the extract was dissolved in glacial acetic acid containing ferric chloride and 1 ml of sulphuric acid was added to the solution. The reaction were observed and recorded.

**Test for alkaloids;** Mayers reagent, Wagners reagent and Drangendoff reagent were added to different test tubes containing the extract solution and each of the reaction was observed and recorded.

#### Experimental Material (*Sorghum bicolor*)

Guinea corn (*Sorghum bicolor*) were obtained from the local market of Samaru Zaria which was cleaned with absolute alcohol after which the seeds were dried before use. The seeds viability was determined by their ability to remain submerged in water. Those that have remained submerged in water were chose and dried for use [8].

#### Determination of growth inhibitory effects of Methanol extracts on guinea corn Seeds radicles length.

About 10ml of 1 – 30 mg/ml of each of the extracts dissolved in 5% dimethyl sulphoxide in water was

poured in to 9cm wide petri dishes laid with cotton wool and filter paper (Whatman no.1). Twenty (20) viable seeds were spread on each plate and incubated in dark environment. The lengths (mm) of radicles emerging from the seeds were taken at 24, 48, 72 and 96 hr. The control seeds were treated with distilled water containing no extracts. The experiment were carried out also in triplicates [8].

#### Statistical Analysis

All data were expressed as mean±SEM and one way Analysis of Variance Anova statistical test using SAS Version 9.2 to test the significance. P<0.05 was considered Significance.

### III. RESULT AND DISCUSSION

#### Preliminary Phytochemical Studies of *Diospyros mespiliformis* Stem Bark

| Test                     | Inference |
|--------------------------|-----------|
| Test for carbohydrate    |           |
| Molisch test             | +         |
| Test for tannins         |           |
| Ferric chloride test     | +         |
| Lead sub-acetate test    | +         |
| Test for saponins        |           |
| Frothing test            | +         |
| Test for sterols         |           |
| Salkowski test           | +         |
| Test for triterpenes     |           |
| Liebermann-burchard test | +         |
| Test for anthraquinones  |           |
| Bontragers test          | +         |
| Test for flavonoids      |           |
| Shinoda test             | +         |
| Sodium hydroxide test    | +         |
| Test for deoxy sugars    |           |
| Keller-kellani test      | +         |
| Test for alkaloids       |           |
| Drangendoff's reagent    | +         |
| Wagners reagent          | +         |
| Mayers reagent           | +         |

**Keywords:** (+) indicates presence of secondary metabolites

#### Thin layer profiling of the Methanol Crude extract



#### Chromatogram of Methanol extract developed in Chloroform:Methanol (4:1)sprayed with Anisaldehyde/ H<sub>2</sub>SO<sub>4</sub>

The 200g of the powdered stem bark of *D. mespiliformis* were observed to have yielded 15.83%

(w/w). The activities of plant extracts in effecting any therapeutic or biological changes in ailing of animals suffering from diseases or living tissues are direct functions of the chemical constituents inherently present in them after extraction [8]. In this research, Methanol extract were tested for the inhibitory effect on the seeds radicles of the Guinea corn (*Sorghum bicolor*). The stem bark of *D. mespiliformis* have been discovered in this work to contain Alkaloids, Flavonoids, Saponins, Steroids and triterpenes, Tannins and Anthraquinones, which are likely to be some of the constituents that contributes to the plants uses in ethnomedicine. Scientific investigation of phytochemicals for probable anti-tumour effects needed allots of materials coupled with the effects and also time even if the result may not be interesting, not showing the expected results of producing more potentially promising materials [9]. These problems have led to the establishment of various bench top methods which have been reportedly used by scientists as indicators of potentially promising anti-tumour phytochemicals e.g [10] and [11] used radicles of germinating seeds. This method is of tremendous value is simple, rapid, reproducible, time and material saving. It is an

experimental procedure which can be carried out in laboratories where appropriate human cells lines are not available. This method can be used to test and evaluate many medicinal plants that may be claimed to treat tumor related diseases. In this research, at 24 hours, the stem bark extract had  $4.771 \pm 0.526$  mm for the controls whereas the seeds treated with 10, 20, and 30 mg/ml of the extract produced a length total of  $2.772 \pm 0.494$  mm,  $2.150 \pm 0.490$  mm and  $2.257 \pm 0.489$  mm respectively. At the end of 96 hours of incubation period, the radicles length of the control seeds measured  $93.77 \pm 9.730$  mm, while those treated with 10, 20, and 30 mg/ml were observed to be  $37 \pm 3.297$  mm,  $17.023 \pm 2.802$  mm and  $16.086 \pm 1.976$  mm respectively as shown in the fig. below. This reduction in the growth implied 60.54, 81.87 and 82.83% respectively compared to the controls as shown in Fig.1 below. This activity might be due to genera relationship between this plant and other species in the same genus as discovered in the *D. undulata* have found to possess the antiproliferative activity against various cancer cells [12]. Another species e.g *D. peregrine* has found to possess antitumor and anti-inflammatory activity [13].




Fig.1: Growth Inhibitory Effect Of The Methanol Extract of the Stem Bark of *D. mespiliformis* on Guinea corn seeds radicles length

**CONCLUSION**

This study has scientifically justified the traditional uses of *Diospyros mespiliformis* stem bark extracts for it has an antiproliferative property against radicles of a Guinea corn (*Sorghum bicolor*) which may relate to its use as anticancer agent. However, use of cancer cell lines will further confirm this claim.

**REFERENCES**

- [1] R.G, Belemtougri. B, Constantin, C, Cognard, G, Raymond, and L, Sawadogo, "Effects of two medicinal plants," *Psidium guajava* L. (Myrtaceae) and *Diospyros mespiliformis* L. (Ebenaceae) leaf extracts on rat skeletal muscle cells in primary culture. *Journal of Zhejiang University*; 7(1): 56-63. Jan. 2006.
- [2] I. E, Mohamed, , E.B, ELNUR, M. I, Choudhary and S.N. Khan, "Bioactive natural products from two Sudanese Medicinal plants *Diospyros mespiliformis* and *Croton zambesicus*," *Resource of Natural Products*.; 3(4): 198-203. Oct. 2009.
- [3] B.S. Adzu, "Pharmacological evidence for curing folkloric use of *D. mespiliformis* Hochst. In the relief of pain and fever", *Journal of Ethnopharmacology* 82, 191-195. Feb. 2002.
- [4] J.B, Harbourne, "*Phytochemical methods*" A guide to modern techniques of plant analysis. Chapman and Hall, London. Pp 49-188. 1973.
- [5] K.R, Brain and T.D, Turner, "*The practical Phytopharmaceutical*". Wright scientecnica Pp 81-144. 1975.
- [6] W.C., Evans, "*Text Book of Pharmacognosy*", SAUNDERS Elsevier Ltd. 15<sup>th</sup> Pp336-7. 2002.
- [7] A . Sofowora , "*Medicinal Plants and Traditional Medicine in Africa*", 3<sup>rd</sup> edn. Spectrum Books Ltd Ibadan-Nigeria, Pp.1-69, 164. 2008.
- [8] B.A, Ayinde. E.K.I, Omogbbai and E.O. Ikpefan, "Comparative cytotoxic and antiproliferative effects of *Persea Americana* mill ( *Lauraceae*) Leaf, stem, and root barks", *Nigerian Journal of Pharmaceutical Sciences*.;10:16-26. March 2011.
- [9] B.A, Ayinde and U. Agbakwuru, "Cytotoxic and growth inhibitory effects of the methanol extract of leaves of *Struchium sparganophora* Ktze (Asteraceae)", *Pharmacognosy magazine*.Vol.6 (24): 293-297. Oct. 2010.
- [10] D. A Shobaike, , A.O, Ogundaini, S.A, Adesanya, "The effects of some synthesized stilbene analogues on *Artemia salina* naupali and germination of *Sorgum bicolor* seeds. *Nigerian Journal of Natural Product and Medicine*; 6:19-25. 2002.
- [11] J.L, McLaughlin, , C. Chang and, D. I, Smith., "*Benchmark biossays for the discovery of bioactive natural products: an update*. In: Atta-ur-Rahman, editor. *Studes in natural products Chemistry*. Vol. 9. Amsterdam: Elsevier Science Publishers; P.383- 409. 1999.
- [12] P.P Rungrojtrakool, J, Siripong , W. Y, Chuakul and R, Temsiririrkul, "Antiproliferative Activity against Various Cancer Cells and Phytochemical Components of Thai Herbal Formula," *Mahidol University Journal of Pharmaceutical Sciences* vol. 39(2), 7-14. 2012.
- [13] V. Y, A, Gopal. G, Ravindranath. A. B, Kalpana and R.V Prabhakar, "Antitumor Activity of *Diospyros peregrina* on Ehrlich Ascites Carcinoma in Mice," *Journal of scientific research*. (2), 413-419. April 2011.

★ ★ ★