

# **The status of ‘on-farm’ seed priming and related work funded by the DFID Plant Sciences Research Programme**

**D Harris**

*Centre for Arid Zone Studies, University of Wales, Bangor, Gwynedd, UK.*

## BACKGROUND

The PSP has always been mindful of the limited resources of our ultimate clients – poor farmers in marginal environments of developing countries. Hence we have pursued a strategy of developing ‘key’ technologies, i.e., simple, low cost, low risk interventions that can be easily adopted, without significant additional capital costs, by farmers. In agriculture, the ideal ‘key’ technology is improved seed. Farmers who adopt new varieties of existing crops can gain immediate benefits without any changes in management, although it is common to find that farmers do increase levels of inputs in response to the reduced risk and additional opportunities afforded by the new varieties. Thus the output of most of the PSP projects is either genetically improved seed (as a result of improved breeding techniques) or a wider choice and better availability of appropriate varieties (by developing participatory varietal selection – PVS – and novel seed supply systems) for resource-poor farmers.

In contrast to the seed based technologies described above, low-cost, low-risk agronomic interventions are less common. Historically, although agronomic improvements in developed countries have contributed substantially to increased crop yields, they have relied on substantial financial investment in machinery, chemical fertilisers and crop protection compounds. Such investments are seldom possible, or even appropriate, for poor farmers in high-risk, rainfed environments. Nevertheless, PSP identified one area – poor crop establishment – where farmers could make improvements without incurring additional costs.

## ON-FARM SEED PRIMING

Harris (1996) highlighted just how difficult it was for farmers in marginal areas to get their crops established effectively but he demonstrated that simply soaking seeds in water before sowing could increase the speed of germination and emergence, leading to better crop stands, and make seedlings grow much more vigorously. These observations were the basis for PSP efforts from 1997 to test, develop and eventually promote seed priming for a wide range of tropical and sub-tropical crops in many developing countries.

### ***Agronomic benefits***

Initially, two projects (R6395 and R7438) used a combination of *in vitro*, on-station and on-farm work to test the germination and emergence characteristics of many crops and varieties in response to seed priming. These two projects had several overall objectives: to test the effects of on-farm seed priming for crops important for resource-poor farmers in marginal areas; to develop, using participatory approaches with farmers, practical methods of priming seeds to best effect; to promote seed priming with resource-poor farmers and organisations (GOs, universities, NGOs) working with such farmers.

The concept of the ‘safe limit’ was found to be important – the maximum length of time for which seeds can be soaked and which, if exceeded, could lead to seed or seedling damage. Once safe limits were determined from *in vitro* studies, participatory work with farmers was used to test and develop through feedback practical methods of priming seeds. Farmers’ trials were also used to assess the effects of priming on a large scale, while further *in vitro* studies

(R7440) and studies on research stations were used to gain insights into possible mechanisms for the effects of priming.

Results quickly began to accumulate that showed that, not only was seed priming effective in improving crop stands (e.g. Fig. 1), but that there were many positive 'knock-on' effects of faster emergence and more vigorous early growth. Priming was seen to promote earlier tillering in cereals, earlier flowering (e.g. Fig. 2) and maturity and higher yields (e.g. Fig. 3). These effects were noted in many crops and countries (see Table 1).



**Figure 1. Better stand and more vigorous growth of chickpea after priming the seeds in Bangladesh.**

**Table 1. Crops in which seed priming has increased yields, the countries involved and the references where the methods used and the results obtained can be found.**

<b>Crop</b>	<b>Countries</b>	<b>References</b>
Wheat	India, Nepal, Pakistan	Harris <i>et al.</i> (2001b); Rashid <i>et al.</i> (2002)
Upland rice	India, Nigeria, Sierra Leone, Gambia, Ghana, Cameroon	Harris <i>et al.</i> (1999); (2002); Harris (2003)
Maize	India, Nepal, Pakistan, Zimbabwe	Harris <i>et al.</i> (1999); (2001a); (2001c)
Sorghum	Pakistan, Botswana, Zimbabwe	Harris (1996); Chivasa <i>et al.</i> (1998); (2001); Rashid <i>et al.</i> (2002)
Pearl millet	Pakistan, India	Harris and Mottram (2004)
Finger millet	India	Kumar <i>et al.</i> (2002)
Chickpea	Bangladesh, India, Nepal, Pakistan	Harris <i>et al.</i> (1999); Musa <i>et al.</i> (2001); Rashid <i>et al.</i> (2002)
Mungbean	Pakistan	Rashid <i>et al.</i> (2004b)
Cowpea	Senegal	Braconnier and Bouru (2004)



**Figure 2. Earlier flowering (left) in maize plants in Zimbabwe from seed primed with water.**

Seed priming is demonstrably effective for a wide range of crops and environments (Table 1), yet its promotion as a stand-alone technology has limitations, particularly for inclusion in government extension programmes. Consequently, more emphasis was placed on using seed priming as one of a set of tools to address more fundamental shortcomings in cropping systems. Other PSP-funded work (R7541) had identified and quantified the huge area of land left fallow in South Asia after the harvest of rainfed rice (Subbarao *et al.*, 2001). Preliminary research (R8098) showed that poor crop establishment was a major constraint on the adoption of a second crop after rice. Projects R7540, R8269 in Bangladesh and R8221 in Eastern India demonstrated that early, rapid, minimum tillage and sowing primed seeds of a short-duration legume such as chickpea was effective in raising yields in these rice fallows (Musa *et al.*, 1999; Musa *et al.*, 2001) to levels where the technology was adopted widely by farmers (Saha, 2002 and this volume).



**Figure 3. Yield of mungbean pods from same-size plots with seeds primed for (L-R) 0, 4, 6 and 8 hours.**

In addition to the acknowledged yield benefits from priming with water alone, PSP research has identified a number of opportunities for priming to be used as a vehicle to introduce fertilising *rhizobia*, additional micronutrients or crop protection agents into seeds.

### **Nutrient supplementation**


Many legumes are relatively unproductive in acid soils because nodulation is limited by poor availability of molybdenum (Mo). In particular, chickpea is known to respond to added Mo but soil application is problematic for resource-poor farmers because of the relatively high rates of application required (at least 0.5 kg/ha) and because uniform application is difficult to achieve. On the other hand, alkaline soils are often deficient in micronutrients such as zinc (Zn). Recent work (R8221 and R8269) in India and Bangladesh has shown that large yield increases in chickpea are possible in farmers' conditions following priming with tiny amounts of molybdenum (Fig. 4; Kumar Rao *et al.*, 2004; Johansen *et al.*, 2004). Similarly, Harris *et al.* (2004b) in R7438 demonstrated large, highly cost-effective yield increases in chickpea and wheat in Pakistan in response to priming with zinc sulphate. Harris *et al.* (2004a) have also shown that seeds can be effectively inoculated with *rhizobium* (to enable fixation of nitrogen) during normal priming operations. Preliminary data (unpublished) also suggest that it is possible for farmers to prime seeds with small amounts of phosphate (P) to good effect in that early root growth is stimulated allowing more effective uptake of available P in the soil. There are huge areas of rice fallows in S. Asia (around 16 million hectares) and many of the soils are deficient in N, P and Mo so the domain for this technology is very large.



**Figure 4. Provision of molybdenum by priming chickpea seed (background) results in better fixation of nitrogen and higher yields in Bangladesh.**

### **Disease resistance**

Musa *et al.* (2001) reported that seed priming in chickpea significantly reduced the damage caused by collar rot (*Sclerotium rolfsii*) in Bangladesh in two contrasting seasons. Recent work in Pakistan (R7438) has demonstrated that mungbean (*Vigna radiata*) grown from seed primed in water for 8 hours before sowing showed significantly fewer serious symptoms of infection by Mungbean Yellow Mosaic Virus (MYMV) than a crop established without priming (Fig. 5). The large differences in virus-related damage were associated with significant increases in pod weight (threefold) and grain weight (fivefold) due to priming (Rashid *et al.*, 2004a). Rashid *et al.* (2004b) also observed similar differences in MYMV infection in other mungbean priming trials.



**Figure 5. Changes in incidence and severity of MYMV disease following seed priming in a mungbean crop in Pakistan.**

These observations from field trials of enhanced resistance to disease following seed priming have prompted an *in vitro* investigation of the phenomenon using a well-established plant-disease model, i.e., pearl millet and downy mildew disease. Ongoing research (R7438) has shown that priming seeds in water for eight hours before sowing significantly reduced the incidence of downy mildew disease in seedlings of a highly susceptible cultivar from about 80% to less than 60% (Table 2 and Harris *et al.*, 2004a).

**Table 2. Percentage of plants showing symptoms of downy mildew disease. Analysis used arcsin transformed data. Transformed means are presented (with non-transformed means in brackets). SE diff. values were calculated using transformed data.**

Treatment	Disease incidence, %.
Seed not primed	63.5 (79.7)
Seed primed for 8 hours	50.3 (58.8)
Standard error of difference between means	3.42
Least significant difference (P<0.05)	7.19

These unexpected results cannot be explained without reference to biochemical changes induced by priming and further experiments are underway (R7438) to investigate this mechanism and to discover if it operates in other plant/disease systems. The practical consequences for resource-poor farmers of such a mechanism are enormous and far-reaching.

### **Promotion and dissemination**

Information on seed priming is available on the website [www.seedpriming.org](http://www.seedpriming.org) and a significant proportion of project staff time is dedicated to the distribution of information related to seed priming. Staff have made presentations on various aspects of seed priming at 20 international conferences and workshops since 1997 and have published widely (Table 1 and reference list below). Almost 3000 copies of the illustrated brochure 'On-Farm' Seed

*Priming. A key technology to improve the livelihoods of resource-poor farmers in marginal environments* have been distributed through various channels. In addition detailed advice, research protocols and supporting materials, e.g. publications concerning seed priming, have been provided in response to enquiries from around 150 individuals and organisations (researchers, NGOs etc) around the world who have expressed an interest in testing seed priming for themselves. Seed priming has been featured on the radio and on TV (twice).

### **The future**

Many thousands of farmers, researchers and extensionists, many not directly linked to PSP, have been exposed to seed priming and enough time has elapsed to allow us to follow up and learn from their experiences. In addition, studies of uptake and persistence of the technology are required where seed priming work has been funded directly by PSP. An example of this is ongoing in Cameroon, Gambia, Ghana, Nigeria and Sierra Leone in the rainfed rice systems described by Harris (2003).

The simplicity and effectiveness of using seed priming to deliver Mo, Zn, P and (via *rhizobium*) N has been demonstrated on a research scale in Bangladesh, India and Pakistan and needs to be researched and disseminated at operational scales. The work so far on second cropping has demonstrated many synergies between priming-related advantages and the introduction of new varieties through PVS – a clear example of the integrated approach favoured by the PSP. The obvious focus for such work is the huge area of rice fallows in South Asia where the livelihoods of some of the world's poorest people can be transformed by growing a low-cost, low-risk second crop after rice. There are also large areas of 'maize fallows' in East Africa and preliminary experiments with chickpea (R7438) have also identified a potential impact there.

Using seed priming to deliver nutrients may need further backup research, particularly for P, in addition to the operational scale, adaptive research noted above. However, more fundamental research will certainly be necessary to understand the nature of the relation between seed priming and disease resistance. Although preliminary research (Musa *et al.*, 2001; Rashid *et al.*, 2004a; Harris *et al.*, 2004b) has established that seed priming can, in some circumstances, increase plant resistance to disease further work is required to confirm this phenomenon to the point where it can be regarded as another of the assured benefits of priming seeds.

## **OUTPUTS FROM SEED PRIMING-RELATED PROJECTS**

*(note: not all references listed below appear in the text)*

2004

Berkelaar, D. (2004). On-farm seed priming. ECHO Development Notes 83: 3-5.

Harris, D. (2004). On-farm seed priming reduces risk and increases yield in tropical crops. In: New Directions for a Diverse Planet. Proceedings of the 4th International Crop Science Congress, 26 Sep - 1 Oct. 2004, Brisbane, Australia.

Harris, D., Breese, W.A. and Kumar Rao, J.V.D.K. (2004a). The improvement of crop yield in marginal environments using 'on-farm' seed priming. In 'Adaptation of Plants to Water-Limited Mediterranean Environments', Floreat, WA, 20-24 September, 2004. CSIRO, Australia.

Harris, D. and Mottram, A. (2004). Practical hydration of seeds of tropical crops: 'on-farm' seed priming. In 'Seed Science and Technology: Trends and Advances', ed. A.S. Basra. The Howarth Press (in press).

Harris, D. Rashid, A. Arif, M. and Yunas, M. (2004b). Alleviating micronutrient deficiencies in alkaline soils of North West Frontier Province of Pakistan: on-farm seed priming with zinc in wheat and chickpea. In 'International Workshop on Agricultural Strategies to Reduce Micronutrient Problems in Mountains and Other Marginal Areas in South and South East Asia'. Kathmandu, 8-10 September, 2004. Nepal Agricultural Research Council.

Harris, D., Rashid, A., Hollington, P.A., Ali, S. and Rafiq, M. (2004c). Mungbean production in the North West Frontier Province of Pakistan and the influence of on-farm seed priming. In 'Improving Income and Nutrition by Incorporating Mungbean in Cereal Fallows in the Indo-Gangetic Plains of South Asia'. Final Workshop and Planning Meeting, Punjab Agricultural University, Ludhiana, Punjab, India, 26-30 May 2004.

Johansen, C., Musa, A.M., Kumar Rao, J.V.D.K., Harris, D., Ali, M.Y. and Lauren, J.G. (2004). Molybdenum response of chickpea in the High Barind Tract of Bangladesh and in Eastern India. In 'International Workshop on Agricultural Strategies to Reduce Micronutrient Problems in Mountains and Other Marginal Areas in South and South East Asia'. Kathmandu, 8-10 September, 2004. Nepal Agricultural Research Council.

Khanal, N., Harris, D., Sherpa, L.T., Giri, R.K. and Joshi, K.D. (2004a). Testing and promotion of mungbean in cereal fallows in the low hills and terai agroecosystems of Nepal. In 'Improving Income and Nutrition by Incorporating Mungbean in Cereal Fallows in the Indo-Gangetic Plains of South Asia'. Final Workshop and Planning Meeting, Punjab Agricultural University, Ludhiana, Punjab, India, 26-30 May 2004.

Khanal, N., Joshi, K. D., Harris, D. and Chand, S. P. (2004b). Effect of micronutrient loading, soil application and foliar sprays of organic extracts on grain legumes and vegetable crops in marginal farmers' condition in Nepal. In 'International Workshop on Agricultural Strategies to Reduce Micronutrient Problems in Mountains and Other Marginal Areas in South and South East Asia'. Kathmandu, 8-10 September, 2004. Nepal Agricultural Research Council.

Kumar Rao, J.V.D.K., Harris, D., Johansen, C. and Musa, A.M. (2004). Low cost provision of molybdenum (Mo) to chickpeas grown in acid soils. Poster in 'International Fertiliser Association Symposium on Micronutrients', New Delhi, 23-25 February, 2004.

Rashid, A., Harris, D., Hollington, P.A. & Ali, S. (2004a). On-farm seed priming reduces yield losses of mungbean (*Vigna radiata*) associated with mungbean yellow mosaic virus in the North West Frontier Province of Pakistan. *Crop Protection* 23: 1119-1124.

Rashid, A., Harris, D., Hollington, P.A. & Rafiq, M. (2004b). Improving the yield of mungbean (*Vigna radiata*) in the North West Frontier Province of Pakistan using on-farm seed priming. *Experimental Agriculture* 40 (2): 233-244.

## 2003

Harris, D. (2003). Reducing risk and increasing yields from rainfed crops in Africa using 'on-farm' seed priming. Pages 87-88 in Abstracts: 'Harnessing Crop Technologies to Alleviate Hunger and Poverty in Africa', 6th Biennial Conference of the African Crop Science Society, Hilton Nairobi Hotel, Kenya, 12th – 16th October, 2003.

Kumar Rao, J.V.D.K., Harris, D., Musa, A.M., Johansen, C., Joshi, K.D., Khanal, N., Gangwar, J.S. and Bhattacharyya, K. (2003). Promotion of rainfed rabi cropping of chickpea in rice fallows of South Asia. Pages 83-84 in Abstracts: International Chickpea Conference (eds: R.N.Sharma, M.Yasin, S.L.Swami, M.A.Khan and Ajit J. William). January 20-22, 2003. Indira Gandhi Agricultural University, Raipur, Chattisgarh, India.

## 2002

Harris, D. (2002a). On-farm seed priming to increase yield of crops and reduce risk of crop failure in marginal areas of developing countries. Pages 1509-1511 in: Second International Agronomy Congress on Balancing Food and Environmental Security – A Continuing Challenge (Extended Summaries) held New Delhi, India, 26-30 November 2002. Indian Society of Agronomy, Indian Council of Agricultural Research and Indian National Academy of Sciences.

Harris, D. (2002b). 'On-farm' seed priming for better crops in marginal areas of developing countries. Pages 128-134 in: Proceedings of the Second International Conference on Sustainable Agriculture for Food, Energy and Industry (Ed. Li Dajue) held 8-13 September 2002, Beijing, China. Institute of Botany, Chinese Academy of Sciences.

Harris, D., Kumar Rao, J.V.D.K. and Kumar, J. (2002a). 'On-farm' seed priming. Pp. 86-88 in: 'Dissemination of technologies enhancing smallholder income in sub-Saharan Africa: Science with a human face.' Proceedings



of an International Workshop held to discuss ICRISAT and World Vision International Partnerships, Myers, R.J.K., Abirifin, A. and Jones, R.B. (eds.) 2002. ICRISAT – Bulawayo, Zimbabwe, 20-23 Nov., 2000.

Harris, D., Rashid, A., Hollington, P.A., Jasi, L. and Riches, C. (2002b). Prospects of improving maize yields with 'on-farm' seed priming. Pp 180-185 in: N.P. Rajbhandari, J.K. Ransom, K. Adikhari and A.F.E. Palmer (eds) 'Sustainable Maize Production Systems for Nepal': Proceedings of a Maize Symposium held, December 3-5, 2001, Kathmandu, Nepal. Kathmandu: NARC and CIMMYT.

Harris, D., Tripathi, R.S. and Joshi, A. (2002c). 'On-farm' seed priming to improve crop establishment and yield in dry direct-seeded rice. Pp. 231-240 in: Pandey, S., Mortimer, M., Wade, L., Tuong, T.P., Lopez, K., and Hardy, B., editors. 2002. Proceedings of the International Workshop on Direct Seeding in Asian Rice Systems: Strategic Research Issues and Opportunities, 25-28 January 2000, Bangkok, Thailand. Los Banos (Philippines): International Rice Research Institute. 383 pp.

Kumar, A., Gangwar, J.S., Prasad, S.C. and Harris, D. (2002). 'On-farm' seed priming increases yield of direct-sown finger millet (*Eleusine coracana*) in India. *International Sorghum and Millets Newsletter* 43: 90-92.

Rashid, A., Harris, D., Hollington, P.A. and Khattak, R.A. (2002). On-farm seed priming: a key technology for improving the livelihoods of resource-poor farmers on saline lands. Pp 423-431 in: R. Ahmad and K.A. Malik (eds.) 'Prospects for Saline Agriculture'. Kluwer Academic Publishers. The Netherlands.

2001

Chivasa, W., Harris, D. and Nyamudeza, P. (2001). On-farm seed priming: a key technology to improve crop establishment and yield in semi-arid tropics. *Sorghum and Millet Improvement Network News* 3 (1): 17-18.

DFID/PSP (2001). 'On-Farm' Seed Priming. A key technology to improve the livelihoods of resource-poor farmers in marginal environments. DFID/PSP information booklet, English language version. Centre for Arid Zone Studies, University of Wales, Bangor, UK.

Harris, D. and Hollington, P.A. (2001). 'On-farm' seed priming – an update. *Tropical Agriculture Association (UK) Newsletter* 21 (4): 7.

Harris, D., Kumar Rao, J.V.D.K. and Kumar, J. (2001a). 'On-farm' seed priming. *Agricultural Research and Extension Network Newsletter* 44: 3.

Harris, D., Pathan, A. K., Gothkar, P., Joshi, A., Chivasa, W. and Nyamudeza, P. (2001b). On-farm seed priming: using participatory methods to revive and refine a key technology. *Agricultural Systems* 69 (1-2): 151-164.

Harris, D., Raghuwanshi, B.S., Gangwar, J.S., Singh, S.C., Joshi, K.D., Rashid, A. and Hollington, P.A. (2001c). Participatory evaluation by farmers of 'on-farm' seed priming in wheat in India, Nepal and Pakistan. *Experimental Agriculture* 37 (3): 403-415.

Musa, A. M., Harris, D., Johansen, C. and Kumar J. (2001). Short duration chickpea to replace fallow after aman rice: the role of on-farm seed priming in the High Barind Tract of Bangladesh. *Experimental Agriculture* 37 (4): 509-521.

Virk, D.S., Harris, D., Raghuwanshi, B.S., Raj, A.G.B, Sodhi, P.S. and Witcombe, J.R. (2001). A holistic approach to participatory crop improvement. Pages 275-282 in 'An Exchange of Experiences from South and South East Asia': Proceedings of the International Symposium on Participatory Plant Breeding and Participatory Plant Genetic Resource Enhancement., held Pokhara, Nepal, 1-5 May 2000. International Center for Tropical Agriculture, Cali, Colombia.

Warham, E and Harris, D. (2001). An old technique – a bucket of water – increases crop yields. *Appropriate Technology* 28 (4) 12-13.

2000

Chivasa, W., Harris, D., Chiduzza, C., Nyamudeza, P. and Mashingaidze, A.B. (2000a).

Biodiversity on-farm in semi-arid agriculture: case study from a smallholder farming system in Zimbabwe. *Zimbabwe Science News* 34: 13-18.

Harris, D. (2000). Participatory testing of 'on-farm' seed priming for direct-seeded rice: a suggested approach for farmer-implemented trials. Pp 17-21 in: 'The Flame Spreads into 2000' Proceedings of the Participatory Rice Improvement and Gender/User Analysis Workshop (PRIGA), held 17-21 April 2000, Bouake, Cote d'Ivoire. West Africa Rice Development Association.

Witcombe, J.R. and Harris, D. (2000). The DFID Plant Sciences Research Programme. *Tropical Agriculture Association (UK) Newsletter* 20 (3): 18-20.

Chivasa, W., Harris, D., Chiduzza, C., Mashingaidze, A.B. and Nyamudeza, P. (2000b). Determination of optimum on-farm seed priming time for maize (*Zea mays* L.) and sorghum (*Sorghum bicolor* [L.] Moench) for

use to improve stand establishment in semi-arid agriculture. *Tanzanian Journal of Agricultural Sciences* 3: No. 2: 103-112.1999

Musa, A.M., Johansen, C., Kumar, J. and Harris, D. (1999). Response of chickpea to seed priming in the High Barind Tract of Bangladesh. *International Chickpea and Pigeonpea Newsletter* 6: 20-22.

Harris, D., Joshi, A., Khan, P.A., Gothkar, P. and Sodhi, P.S. (1999). On-farm seed priming in semi-arid agriculture: development and evaluation in maize, rice and chickpea in India using participatory methods. *Experimental Agriculture* 35: 15-29.

1998

Chivasa, W., Harris, D., Chiduza, C., Nyamudeza, P. and Mashingaidze, A.B. (1998). Agronomic practices, major crops and farmers' perceptions of the importance of good stand establishment in Musikavanhu Communal Area, Zimbabwe. *Journal of Applied Sciences in Southern Africa* 4 (2): 9-25.

Harris, D., Khan, P.A., Gothkar, P., Joshi, A., Raguwanshi, B.S., Parey, A., Sodhi, P.S., Virk, D.S. and Witcombe, J.R. (1998). Key technologies: more appropriate varieties and on-farm seed priming. *Tropical Agriculture Association (UK) Newsletter* 18 (2): 51-53.

Howarth, C.J., Weltzien Rattunde, E., Bidinger, F.R. and Harris, D. (1998). Seedling survival of abiotic stress: sorghum and pearl millet. In *Proceedings of the International Conference on Genetic Improvement of Sorghum and Pearl Millet*, ICRISAT/INTSORMIL, Lubbock, Texas, 22-27 September, 1996. pp 379-399.

1997

Harris, D. & Jones, M (1997). On-farm seed priming to accelerate germination in rainfed, dry-seeded rice. *International Rice Research Notes* 22 (2) 30.

1996

Harris, D. (1996). The effects of manure, genotype, seed priming, depth and date of sowing on the emergence and early growth of *Sorghum bicolor* (L.) Moench in semi-arid Botswana. *Soil & Tillage Research* 40 73-88.

#### Other Outputs

Harris, D. (2002). Getting started quickly: crop establishment and 'on-farm' seed priming. Invited lecture, Cornell University, 29th January 2002. Ithaca, New York, USA.

Harris, D., Khan, P.A., Gothkar, P., Joshi, A., Raguwanshi, B.S., Parey, A. & Sodhi, P.S. (1998). Using participatory methods to develop, test and promote on-farm seed priming in India. In 'International Conference on Food Security and Crop Science', November 3-6, 1998, Hisar, India.

Harris, D., Khan, P.A., Gothkar, P., Joshi, A., Raguwanshi, B.S., Parey, A., Sodhi, P.S., Virk, D.S. and Witcombe, J.R. (1998). Key technologies: more appropriate varieties and on-farm seed priming. Invited presentation at meeting of Tropical Agriculture Association, Long Ashton, 26 March, 1998.

Witcombe, J.R. and Harris, D. (1997). Impact of the DFID Plant Science Research Programme. Paper presented at the DFID Natural Resources Advisers' Conference, Sparsholt, U.K. 6-11 June, 1997. Discussion paper no. 3. Centre for Arid Zone Studies, University of Wales.