

Rice

Rice is the seed of the monocot plants *Oryza sativa* (Asian rice) or *Oryza glaberrima* (African rice). As a cereal grain, it is the most widely consumed staple food for a large part of the world's human population, especially in Asia and the West Indies. It is the grain with the second-highest worldwide production, after maize (corn), according to data for 2010.^[1]

Since a large portion of maize crops are grown for purposes other than human consumption, rice is the most important grain with regard to human nutrition and caloric intake, providing more than one fifth of the calories consumed worldwide by the human species.^[2]

There are many varieties of rice and culinary preferences tend to vary regionally. In the Far East, there is a preference for softer and stickier varieties. Because of its importance as a staple food, rice has considerable cultural importance. Rice is often directly associated with prosperity and fertility. Therefore, there is the custom of throwing rice at weddings.^[3]



Oryza sativa, commonly known as Asian rice

Rice is normally grown as an annual plant, although in tropical areas it can survive as a perennial and can produce a ratoon crop for up to 30 years.^[4] The rice plant can grow to 1–1.8 m (3.3–5.9 ft) tall, occasionally more depending on the variety and soil fertility. It has long, slender leaves 50–100 cm (20–39 in) long and 2–2.5 cm (0.79–0.98 in) broad. The small wind-pollinated flowers are produced in a branched arching to pendulous inflorescence 30–50 cm (12–20 in) long. The edible seed is a grain (caryopsis) 5–12 mm (0.2–0.47 in) long and 2–3 mm (0.079–0.12 in) thick.

Rice cultivation is well-suited to countries and regions with low labor costs and high rainfall, as it is labor-intensive to cultivate and requires ample water. Rice can be grown practically anywhere, even on a steep hill or mountain. Although its parent species are native to Asia and certain parts of Africa, centuries of trade and exportation have made it commonplace in many cultures worldwide.

The traditional method for cultivating rice is flooding the fields while, or after, setting the young seedlings. This simple method requires sound planning and servicing of the water damming and channeling, but reduces the growth of less robust weed and pest plants that have no submerged growth state, and deters vermin. While flooding is not mandatory for the cultivation of rice, all other methods of irrigation require higher effort in weed and pest control during growth periods and a different approach for fertilizing the soil.

The name **wild rice** is usually used for species of the grass genus *Zizania*, both wild and domesticated, although the term may also be used for primitive or uncultivated varieties of *Oryza*.

Etymology

First attested in English in the middle of the 13th century, the word "rice" derives from the Old French *ris*, which comes from Italian *riso*, in turn from the Latin *oriza*, which derives from the Greek ὄρυζα (*oruzá*). The Greek word is the source of all European words (cf. Welsh *reis*, Ger. *Reis*, Lith. *ryžiai*, Serbo-Cr. *riža*, Pol. *ryż*, Dutch *rijst*, Romanian *orez*).^{[5][6][7]} The origin of the Greek word is unclear. It has sometimes held to be from the Tamil word அரிசி (*arisi*), or rather Old Tamil *arici*.^{[8][9]} However, Krishnamurti^[10] disagrees with the notion that Old Tamil *arici* is the source of the Greek term, and proposes that it was borrowed from descendants of Proto-Dravidian **wariñci* instead. Mayrhofer^[11] suggests that the immediate source of the Greek word is to be sought in Old Iranian

words of the types **vrīz-* or **vrinj-*, but these are ultimately traced back to Indo-Aryan (as in Sanskrit *vrīhí-*) and subsequently to Dravidian by Witzel and others.

Preparation as food



Rice broker in 1820s Japan ("*36 Views of Mount Fuji*" Hokusai)

The seeds of the rice plant are first milled using a rice huller to remove the chaff (the outer husks of the grain). At this point in the process, the product is called brown rice. The milling may be continued, removing the bran, *i.e.*, the rest of the husk and the germ, thereby creating white rice. White rice, which keeps longer, lacks some important nutrients; moreover, in a limited diet which does not supplement the rice, brown rice helps to prevent the disease beriberi.

Either by hand or in a rice polisher, white rice may be buffed with glucose or talc powder (often called polished rice, though this term may also refer to white rice in general), parboiled, or processed into flour. White rice may also be enriched by adding nutrients, especially those lost during the milling process. While the cheapest method of enriching involves adding a powdered blend of nutrients that will easily wash off (in the United States, rice which has been so treated requires a label warning against rinsing), more sophisticated methods apply nutrients directly to the grain, coating the grain with a water-insoluble substance which is resistant to washing.



Thresh rice cleaning by a watermill ("*36 Views of Mount Fuji*" Hokusai)

In some countries, a popular form, parboiled rice, is subjected to a steaming or parboiling process while still a brown rice grain. This causes nutrients from the outer husk, especially thiamine, to move into the grain itself. The parboil process causes a gelatinisation of the starch in the grains. The grains become less brittle, and the color of the milled grain changes from white to yellow. The rice is then dried, and can then be milled as usual or used as brown rice. Milled parboiled rice is nutritionally superior to standard milled rice. Parboiled rice has an additional benefit in that it does not stick to the pan during cooking, as happens when cooking regular white rice. This type of rice is eaten in parts of India and countries of West Africa are also accustomed to consuming parboiled rice.



Transplanting rice seedlings in Java, Indonesia

Despite the hypothetical health risks of talc (such as stomach cancer),^[12] talc-coated rice remains the norm in some countries due to its attractive shiny appearance, but it has been banned in some, and is no longer widely used in others (such as the United States). Even where talc is not used, glucose, starch, or other coatings may be used to improve the appearance of the grains.



Removing seedlings from seed beds, the first step along a labor-intensive process of growing rice as seen in Cambodia



Planting rice (San Nicolas, Philippines).

Rice bran, called *nuka* in Japan, is a valuable commodity in Asia and is used for many daily needs. It is a moist, oily inner layer which is heated to produce oil. It is also used as a pickling bed in making rice bran pickles and *takuan*.

Raw rice may be ground into flour for many uses, including making many kinds of beverages, such as *amazake*, *horchata*, rice milk, and rice wine. Rice flour does not contain gluten, so is suitable for people on a gluten-free diet. Rice may also be made into various types of noodles. Raw, wild, or brown rice may also be consumed by raw-foodist or fruitarians if soaked and sprouted (usually a week to 30 days – gaba rice).

Processed rice seeds must be boiled or steamed before eating. Boiled rice may be further fried in cooking oil or butter (known as fried rice), or beaten in a tub to make *mochi*.

Rice, white, long-grain vegetable, raw

Nutritional value per 100 g (3.5 oz)	
Energy	1,527 kJ (365 kcal)
Carbohydrates	80 g
- Sugars	0.12 g
- Dietary fiber	1.3 g
Fat	0.66 g
Protein	7.13 g
Water	11.61 g
Thiamine (vit. B ₁)	0.0701 mg (6%)
Riboflavin (vit. B ₂)	0.0149 mg (1%)
Niacin (vit. B ₃)	1.62 mg (11%)
Pantothenic acid (B ₅)	1.014 mg (20%)
Vitamin B ₆	0.164 mg (13%)
Calcium	28 mg (3%)
Iron	0.80 mg (6%)
Magnesium	25 mg (7%)
Manganese	1.088 mg (52%)
Phosphorus	115 mg (16%)
Potassium	115 mg (2%)
Zinc	1.09 mg (11%)
Percentages are relative to US recommendations for adults. Source: USDA Nutrient Database ^[13]	

Rice is a good source of protein and a staple food in many parts of the world, but it is not a complete protein: it does not contain all of the essential amino acids in sufficient amounts for good health, and should be combined with other sources of protein, such as nuts, seeds, beans, fish, or meat.^[14]

Rice, like other cereal grains, can be puffed (or popped). This process takes advantage of the grains' water content and typically involves heating grains in a special chamber. Further puffing is sometimes accomplished by processing puffed pellets in a low-pressure chamber. The ideal gas law means either lowering the local pressure or raising the water temperature results in an increase in volume prior to water evaporation, resulting in a puffy texture. Bulk raw rice density is about 0.9 g/cm³. It decreases to less than one-tenth that when puffed.

Cooking

The many varieties of rice, for many purposes, are distinguished as long-, medium-, and short-grain rices. The grains of fragrant long-grain rice (high amylose) tend to remain intact after cooking; medium-grain rice (high amylopectin) becomes more sticky. Medium-grain rice is used for sweet dishes, for *risotto* in Italy and many rice dishes, such as *arròs negre*, in Spain. Some varieties of long-grain rice are high in amylopectin, these are generally known as Thai Sticky rice, usually steamed.^[15] A stickier medium-grain rice is used for *sushi*; the stickiness lets the rice be moulded into a solid shape. Short-grain rice is often used for rice pudding.

Rice is cooked by boiling or steaming, and absorbs water during cooking. It can be cooked in just as much water as it absorbs (the absorption method), or in a large quantity of water which is drained before serving (the rapid-boil method).^[16] Electric rice cookers, popular in Asia and Latin America, simplify the process of cooking rice. Rice (or any other grain) is sometimes quickly fried in oil or fat before boiling (for example saffron rice or risotto); this makes the cooked rice less sticky, and is a cooking style commonly called pilaf by American chefs or biryani (Dam-pukhtak) in India, Pakistan, and Iran.



Traditional Medicinal Rice of Niyamgiri Hills,
India



Uncooked, polished, white long-grain rice grains



A Pakistani-style Chinese rice dish using the
Basmati variety



Rice served with Indian curry – note the yellowish tinge in the rice, due to the common Indian practice of adding turmeric during cooking.



Unmilled to milled rice, from left to right, white rice (*Japanese rice*), rice with germ, brown rice




Black rice – one of several colored varieties of natural rice, rich in anthocyanins and tocopherols. Image compares white and black rice from Japan.

In Arab cuisine, rice is an ingredient of many soups and dishes with fish, poultry, and other types of meat. It is also used to stuff vegetables or is wrapped in grape leaves (dolma). When combined with milk, sugar, and honey, it is used to make desserts. In some regions, such as Tabaristan, bread is made using rice flour. Medieval Islamic texts spoke of medical uses for the plant.^[17]

Rice may also be made into congee (also called rice porridge, fawrclaab, okayu, Xifan, jook, or rice gruel) by adding more water than usual, so that the cooked rice is saturated with water, usually to the point that it disintegrates. Rice porridge is commonly eaten as a breakfast food, and is also a traditional food for the sick.



Rice stem cross section magnified 400 times



Rice may be soaked prior to cooking, which saves fuel, decreases cooking time, minimizes exposure to high temperature and thus decreases the stickiness of the rice. For some varieties, soaking improves the texture of the cooked rice by increasing expansion of the grains.

Instant rice differs from parboiled rice in that it is milled, fully cooked and then dried. There is a significant degradation in taste and texture.

A nutritionally superior method of preparing brown rice known as **GABA Rice** or GBR (Germinated Brown Rice)^[18] may be used. This involves soaking washed brown rice for 20 hours in warm water (38 °C or 100 °F) prior to cooking it. This stimulates germination, which activates various enzymes in the rice. By this method, a result of research carried out for the United Nations International Year of Rice, it is possible to obtain a more complete amino acid profile, including GABA.

Rice flour and starch often are used in batters and breadings to increase crispiness.

Nutrition and Health

Nutrients and the nutritional importance of rice

Rice is the staple food of over half the world's population. It is the predominant dietary energy source for 17 countries in Asia and the Pacific, 9 countries in North and South America and 8 countries in Africa. Rice provides 20% of the world's dietary energy supply, while wheat supplies 19% and maize 5%.^[19]

A detailed analysis of nutrient content of rice suggests that the nutrition value of rice varies based on a number of factors. It depends on the strain of rice, that is between white, brown, black, red and purple varieties of rice – each prevalent in different parts of the world. It also depends on nutrient quality of the soil rice is grown in, whether and how the rice is polished or processed, the manner it is enriched, and how it is prepared before consumption.^[20]

An illustrative comparison between white and brown rice of protein quality, mineral and vitamin quality, carbohydrate and fat quality suggests that neither is a complete nutrition source. Between the two, there is a significant difference in fiber content and minor differences in other nutrients.^[21]



Brilliantly colored rice strains such as the purple rice derives its color from anthocyanins and tocopherols. Scientific studies suggest that these color pigments have antioxidant properties that may be useful to human health. In purple rice bran, hydrophilic antioxidants are in greater quantity and have higher free radical scavenging activity than lipophilic antioxidants. Anthocyanins and γ -tocopherols in purple rice are largely located in the inner portion of purple rice bran.^[22]

Comparative nutrition studies on red, black and white varieties of rice suggest that pigments in red and black rice varieties may offer nutrition benefits. Red or black rice consumption was found to reduce or retard the progression of atherosclerotic plaque development, induced by dietary cholesterol, in mammals. White rice consumption offered no similar benefits, and the study claims this to be due to absent antioxidants in red and black varieties of rice.^[23]

Comparison of rice to other major staple foods

The table below shows the nutrient content of major staple foods in a raw form. Raw grains, however, aren't edible and can not be digested. These must be sprouted, or prepared and cooked for human consumption. In sprouted and cooked form, the relative nutritional and anti-nutritional contents of each of these grains is remarkably different from that of raw form of these grains reported in this table.

Nutrient content of major staple foods^[24]

STAPLE:	Maize / Corn ^[A]	Rice ^[B]	Wheat ^[C]	Potato ^[D]	Cassava ^[E]	Soybean ^[F]	Sweet potato ^[G]	Sorghum ^[H]	Yam ^[Y]	Plantain ^[Z]
Component (per 100g portion)	Amount	Amount	Amount	Amount	Amount	Amount	Amount	Amount	Amount	Amount
Water (g)	76	12	11	79	60	68	77	9	70	65
Energy (kJ)	360	1528	1419	322	670	615	360	1419	494	511
Protein (g)	3.2	7.1	13.7	2.0	1.4	13.0	1.6	11.3	1.5	1.3
Fat (g)	1.18	0.66	2.47	0.09	0.28	6.8	0.05	3.3	0.17	0.37
Carbohydrates (g)	19	80	71	17	38	11	20	75	28	32
Fiber (g)	2.7	1.3	10.7	2.2	1.8	4.2	3	6.3	4.1	2.3
Sugar (g)	3.22	0.12	0	0.78	1.7	0	4.18	0	0.5	15
Calcium (mg)	2	28	34	12	16	197	30	28	17	3
Iron (mg)	0.52	4.31	3.52	0.78	0.27	3.55	0.61	4.4	0.54	0.6
Magnesium (mg)	37	25	144	23	21	65	25	0	21	37
Phosphorus (mg)	89	115	508	57	27	194	47	287	55	34
Potassium (mg)	270	115	431	421	271	620	337	350	816	499
Sodium (mg)	15	5	2	6	14	15	55	6	9	4
Zinc (mg)	0.45	1.09	4.16	0.29	0.34	0.99	0.3	0	0.24	0.14
Copper (mg)	0.05	0.22	0.55	0.11	0.10	0.13	0.15	-	0.18	0.08
Manganese (mg)	0.16	1.09	3.01	0.15	0.38	0.55	0.26	-	0.40	-
Selenium (mcg)	0.6	15.1	89.4	0.3	0.7	1.5	0.6	0	0.7	1.5
Vitamin C (mg)	6.8	0	0	19.7	20.6	29	2.4	0	17.1	18.4
Thiamin (mg)	0.20	0.58	0.42	0.08	0.09	0.44	0.08	0.24	0.11	0.05
Riboflavin (mg)	0.06	0.05	0.12	0.03	0.05	0.18	0.06	0.14	0.03	0.05
Niacin (mg)	1.70	4.19	6.74	1.05	0.85	1.65	0.56	2.93	0.55	0.69

Pantothenic acid (mg)	0.76	1.01	0.94	0.30	0.11	0.15	0.80	-	0.31	0.26
Vitamin B6 (mg)	0.06	0.16	0.42	0.30	0.09	0.07	0.21	-	0.29	0.30
Folate Total (mcg)	46	231	43	16	27	165	11	0	23	22
Vitamin A (IU)	208	0	0	2	13	180	14187	0	138	1127
Vitamin E, alpha-tocopherol (mg)	0.07	0.11	0	0.01	0.19	0	0.26	0	0.39	0.14
Vitamin K (mcg)	0.3	0.1	0	1.9	1.9	0	1.8	0	2.6	0.7
Beta-carotene (mcg)	52	0	0	1	8	0	8509	0	83	457
Lutein+zeaxanthin (mcg)	764	0	0	8	0	0	0	0	0	30
Saturated fatty acids (g)	0.18	0.18	0.45	0.03	0.07	0.79	0.02	0.46	0.04	0.14
Monounsaturated fatty acids (g)	0.35	0.21	0.34	0.00	0.08	1.28	0.00	0.99	0.01	0.03
Polyunsaturated fatty acids (g)	0.56	0.18	0.98	0.04	0.05	3.20	0.01	1.37	0.08	0.07

- | | |
|--------------------------------|---|
| A corn, sweet, yellow, raw | B rice, white, long-grain, regular, raw |
| C wheat, durum | D potato, flesh and skin, raw |
| E cassava, raw | F soybeans, green, raw |
| G sweetpotato, raw, unprepared | H sorghum, raw |
| Y yam, raw | Z plantains, raw |

Arsenic concerns

Rice and rice products contain arsenic, a known poison and Group 1 carcinogen.^[25] There is no safe level of arsenic, but, as of 2012, a limit of 10 parts per billion has been established in the United States for drinking water, twice the level of 5 parts per billion originally proposed by the EPA. Consumption of one serving of some varieties of rice gives more exposure to arsenic than consumption of 1 liter of water that contains 5 parts per billion arsenic; however, the amount of arsenic in rice varies widely with the greatest concentration in brown rice and rice grown on land formerly used to grow cotton; in the United States, Arkansas, Louisiana, Missouri, and Texas.^[26] The US FDA is currently studying this issue, but has not established a limit.^[27] China has set a limit of 150 ppb for arsenic in rice^[28].

White rice grown in Arkansas, Louisiana, Missouri, and Texas, which account for 76 percent of American-produced rice, had higher levels of arsenic, possibly because of past use of arsenic based pesticides to control cotton weevils.^[29] Rice from Thailand and India contain the least arsenic among rice varieties in one study.^[30]

Bacillus cereus

Cooked rice can contain *Bacillus cereus* spores, which produce an emetic toxin when left at 4–60 °C (39–140 °F). When storing cooked rice for use the next day, rapid cooling is advised to reduce the risk of toxin production.^[31] One of the enterotoxins produced by *Bacillus cereus* is heat-resistant; reheating contaminated rice kills the bacteria, but does not destroy the toxin already present.

Rice growing ecology

Rice can be grown in different environments, depending upon water availability.^[32] Generally, rice does not thrive in a waterlogged area, yet it can survive and grow herein^[33] and it can also survive flooding.^[34]


1. **Lowland, rainfed**, which is drought prone, favors medium depth; waterlogged, submergence, and flood prone
2. **Lowland, irrigated**, grown in both the wet season and the dry season
3. **Deep water** or floating rice
4. **Coastal Wetland**
5. **Upland rice** is also known as '**Ghaiya rice**', well known for its drought tolerance^[35]

History of domestication and cultivation

Asia

There have been plenty of debates on the origins of the domesticated rice. However, in 2011, a combined effort by Stanford University, New York University, Washington University, and Purdue University has provided DNA evidence that all forms of Asian rice spring from a single domestication of the wild rice *Oryza rufipogon*.^[36] A 2012 study indicated the domestication occurred in the Pearl River valley of China.^[37] Before this research, the commonly accepted view is that rice was first domesticated in the region of the Yangtze River valley in China.^{[38][39]} Morphological studies of rice phytoliths from the Diaotonghuan archaeological site clearly show the transition from the collection of wild rice to the cultivation of domesticated rice. The large number of wild rice phytoliths at the Diaotonghuan level dating from 12,000–11,000 BP indicates that wild rice collection was part of the local means of subsistence. Changes in the morphology of Diaotonghuan phytoliths dating from 10,000–8,000 BP show that rice had by this time been domesticated.^[40] Soon afterwards the two major varieties of indica and japonica rice were being grown in Central China.^[39] In the late 3rd millennium BC, there was a rapid expansion of rice cultivation into mainland Southeast Asia and westwards across India and Nepal.^[39]

In 2003, Korean archaeologists claimed to have discovered the world's oldest domesticated rice.^[41] Their 15,000-year old age challenges the accepted view that rice cultivation originated in China about 12,000 years ago.^[41] These findings were received by academia with strong skepticism,^[42] and the results and their publicizing has been cited as



being driven by a combination of nationalist and regional interests.^[43] In 2011, a combined effort by the Stanford University, New York University, Washington University in St. Louis, and Purdue University has provided the strongest evidence yet that there is only one single origin of domesticated rice, in the Yangtze Valley of China.^{[44][45]}

The earliest remains of the grain in the Indian subcontinent have been found in the Indo-Gangetic Plain and date from 7000–6000 BC though the earliest widely accepted date for cultivated rice is placed at around 3000–2500 BC with findings in regions belonging to the Indus Valley Civilization. Perennial wild rices still grow in Assam and Nepal. It seems to have appeared around 1400 BC in southern India after its domestication in the northern plains. It then spread to all the fertile alluvial plains watered by rivers. Cultivation and cooking methods are thought to have spread to the west rapidly and by medieval times, southern Europe saw the introduction of rice as a hearty grain.

According to Zohary and Hopf (2000, p. 91), *O. sativa* was recovered from a grave at Susa in Iran (dated to the 1st century AD) at one end of the ancient world, another domestication of rice in South Asia.

Today, the majority of all rice produced comes from China, India, Indonesia, Bangladesh, Vietnam, Thailand, Myanmar, Pakistan, Philippines, and Japan. Asian farmers still account for 92% of the world's total rice production.

Companion plant

One of the earliest known examples of companion planting is the growing of rice with Azolla, the mosquito fern, which covers the top of a fresh rice paddy's water, blocking out any competing plants, as well as fixing nitrogen from the atmosphere for the rice to use. The rice is planted when it is tall enough to poke out above the azolla. This method has been used for at least a thousand years.

Africa

African rice has been cultivated for 3500 years. Between 1500 and 800 BC, *Oryza glaberrima* propagated from its original centre, the Niger River delta, and extended to Senegal. However, it never developed far from its original region. Its cultivation even declined in favour of the Asian species, which was introduced to East Africa early in the common era and spread westward.^[46] African rice helped Africa conquer its famine of 1203.^[47]

Middle East



Semi-ripe Paddy in West Bengal, India



The Banaue Rice Terraces in the Philippines.



Rice field under monsoon clouds in Pegu Division, Burma



Rice crop in Madagascar

Rice was grown in some areas of southern Iraq. With the rise of Islam it moved north to Nisibin, the southern shores of the Caspian Sea (Iran) and then beyond the Muslim world into the valley of Volga. In Egypt, rice is mainly grown in the Nile Delta. In Palestine, rice came to be grown in the Jordan Valley. Rice is also grown in Saudi Arabia at Al-Hasa Oasis and Yemen.^[48]

Europe

Rice was known to the Classical world, being imported from Egypt, and perhaps west Asia. It was known to Greece by returning soldiers from Alexander III's ("the Great") military expedition to Asia. Large deposits of rice from the first century A.D. have been found in Roman camps in Germany.^[49]

The Moors brought Asiatic rice to the Iberian Peninsula in the 10th century. Records indicate it was grown in Valencia and Majorca. In Majorca, rice cultivation seems to have stopped after the Christian conquest, although historians are not certain.^[48]

Muslims also brought rice to Sicily, where it was an important crop^[48] long before it is noted in the plain of Pisa (1468) or in the Lombard plain (1475), where its cultivation was promoted by Ludovico Sforza, Duke of Milan, and demonstrated in his model farms.^[50]

After the 15th century, rice spread throughout Italy and then France, later propagating to all the continents during the age of European exploration.

Caribbean and Latin America

Rice is not native to the Americas but was introduced to Latin America and the Caribbean by European colonizers at an early date with Spanish colonizers introducing Asian rice to Mexico in the 1520s at Veracruz and the Portuguese and their African slaves introducing it at about the same time to Colonial Brazil.^[51] Recent scholarship suggests that enslaved Africans played an active role in the establishment of rice in the New World and that African rice was an important crop from an early period.^[52] Varieties of rice and bean dishes that were a staple dish along the peoples of West Africa remained a staple among their descendants subjected to slavery in the Spanish New World colonies, Brazil and elsewhere in the Americas.^[47]

The Native Americans of what is now the Eastern United States may have practiced extensive agriculture with forms of wild rice.

United States

In 1694, rice arrived in South Carolina, probably originating from Madagascar.^[51]

In the United States, colonial South Carolina and Georgia grew and amassed great wealth from the Slavery labor obtained from the Senegambia area of West Africa and from coastal Sierra Leone. At the port of Charleston, through which 40% of all American slave imports passed, slaves from this region of Africa brought the highest prices, in recognition of their prior knowledge of rice culture, which was put to use on the many rice plantations around Georgetown, Charleston, and Savannah. From the enslaved Africans, plantation owners learned how to dyke the marshes and periodically flood the fields. At first the rice was milled by hand with wooden paddles, then winnowed in sweetgrass baskets (the making of which was another skill brought by slaves from Africa). The invention of the rice mill increased profitability of the crop, and the addition of water power for the mills in 1787 by millwright Jonathan Lucas was another step forward. Rice culture in the southeastern U.S. became less profitable with the loss of slave



South Carolina rice plantation, showing a Winnowing barn (Mansfield Plantation, Georgetown)

labor after the American Civil War, and it finally died out just after the turn of the 20th century. Today, people can visit the only remaining rice plantation in South Carolina that still has the original winnowing barn and rice mill from the mid-19th century at the historic Mansfield Plantation in Georgetown, South Carolina. The predominant strain of rice in the Carolinas was from Africa and was known as "Carolina Gold." The cultivar has been preserved and there are current attempts to reintroduce it as a commercially grown crop.^[53]

In the southern United States, rice has been grown in southern Arkansas, Louisiana, and east Texas since the mid-19th century. Many Cajun farmers grew rice in wet marshes and low lying prairies where they could also farm crayfish when the fields were flooded.^[54] In recent years rice production has risen in North America, especially in the Mississippi River Delta areas in the states of Arkansas and Mississippi.

Rice cultivation began in California during the California Gold Rush, when an estimated 40,000 Chinese laborers immigrated to the state and grew small amounts of the grain for their own consumption. However, commercial production began only in 1912 in the town of Richvale in Butte County.^[55] By 2006, California produced the second largest rice crop in the United States,^[56] after Arkansas, with production concentrated in six counties north of Sacramento.^[57] Unlike the Mississippi Delta region, California's production is dominated by short- and medium-grain *japonica* varieties, including cultivars developed for the local climate such as Calrose, which makes up as much as 85% of the state's crop.^[58]

References to wild rice in the Americas are to the unrelated *Zizania palustris*

More than 100 varieties of rice are commercially produced primarily in six states (Arkansas, Texas, Louisiana, Mississippi, Missouri, and California) in the U.S.^[59] According to estimates for the 2006 crop year, rice production in the U.S. is valued at \$1.88 billion, approximately half of which is expected to be exported. The U.S. provides about 12% of world rice trade.^[59] The majority of domestic utilization of U.S. rice is direct food use (58%), while 16% is used in each of processed foods and beer. The remaining 10% is found in pet food.^[59]

Australia

Rice was one of the earliest crops planted in Australia by British settlers, who had experience with rice plantations in the Americas and the subcontinent.

Although attempts to grow rice in the well-watered north of Australia have been made for many years, they have consistently failed because of inherent iron and manganese toxicities in the soils and destruction by pests.

In the 1920s it was seen as a possible irrigation crop on soils within the Murray-Darling Basin that were too heavy for the cultivation of fruit and too infertile for wheat.^[60]

Because irrigation water, despite the extremely low runoff of temperate Australia, was (and remains) very cheap, the growing of rice was taken up by agricultural groups over the following decades. Californian varieties of rice were found suitable for the climate in the Riverina, and the first mill opened at Leeton in 1951.

Even before this Australia's rice production greatly exceeded local needs,^[60] and rice exports to Japan have become a major source of foreign currency. Above-average rainfall from the 1950s to the middle 1990s^[61] encouraged the expansion of the Riverina rice industry, but its prodigious water use in a practically waterless region began to attract the attention of environmental scientists. These became severely concerned with declining flow in the Snowy River and the lower Murray River.


Although rice growing in Australia is highly profitable due to the cheapness of land, several recent years of severe drought have led many to call for its elimination because of its effects on extremely fragile aquatic ecosystems. The Australian rice industry is somewhat opportunistic, with the area planted varying significantly from season to season depending on water allocations in the Murray and Murrumbidgee irrigation regions.

Production and commerce

Top 20 Rice Producers by Country — 2010 (million metric ton) ^[62]	
 People's Republic of China	197.2
 India	120.6
 Indonesia	66.4
 Bangladesh	49.3
 Vietnam	39.9
 Burma	33.2
 Thailand	31.5
 Philippines	15.7
 Brazil	11.3
 United States	11.0
 Japan	10.6
 Cambodia	8.2
 Pakistan	7.2
 South Korea	6.1
 Madagascar	4.7
 Egypt	4.3
 Sri Lanka	4.3
 Nepal	4.0
 Nigeria	3.2
 Laos	3.0

Source: Food and Agriculture Organization

Production



Rice is a major food staple and a mainstay for the rural population and their food security. It is mainly cultivated by small farmers in holdings of less than 1 hectare. Rice is also a wage commodity for workers in the cash crop or non-agricultural sectors. Rice is vital for the nutrition of much of the population in Asia, as well as in Latin America and the Caribbean and in Africa; it is central to the food security of over half the world population. Developing countries account for 95% of the total production, with China and India alone responsible for nearly half of the world output.^[63]

World production of rice has risen steadily from about 200 million tonnes of paddy rice in 1960 to over 678 million tonnes in 2009. The three largest producers of rice in 2009 were China (197 million tonnes), India (131 Mt), and Indonesia (64 Mt). Among the six largest rice producers, the most productive farms for rice, in 2009, were in China producing 6.59 tonnes per hectare.^[64] At 44 million hectares, India had the largest farm area under rice production in 2009. The rice farm productivity in India were about 45% of the rice farm productivity in China, and about 60% of

the rice farm productivity in Indonesia. If India could adopt the farming knowledge and technology in use in China and Indonesia, India could produce an additional 100 million tonnes of rice, enough staple food for about 400 million people every year, and US\$50 billion in additional annual income to its rice farmers (adjusted to 2010 dollars and global rice prices per tonne). In the 1990s, genetic studies took place in many European laboratories to increase rice production per hectare. Most of them were Dutch agricultural organizations united by HNGAC. These studies were later stopped due to lack of funding.

In addition to the gap in farming system technology and knowledge, many rice grain producing countries have significant losses post-harvest at the farm and because of poor roads, inadequate storage technologies, inefficient supply chains and farmer's inability to bring the produce into retail markets dominated by small shopkeepers. A World Bank – FAO study claims 8% to 26% of rice is lost in developing nations, on average, every year, because of post-harvest problems and poor infrastructure. Some sources claim the post-harvest losses to exceed 40%.^{[63][65]} Not only do these losses reduce food security in the world, the study claims that farmers in developing countries such as China, India and others lose approximately US\$89 billion of income in preventable post-harvest farm losses, poor transport, the lack of proper storage and retail. One study claims that if these post-harvest grain losses could be eliminated with better infrastructure and retail network, in India alone enough food would be saved every year to feed 70 to 100 million people over a year.^[66]

Harvesting, drying and milling

Unmilled rice, known as paddy (Indonesia and Malaysia: padi; Philippines, palay), is usually harvested when the grains have a moisture content of around 25%. In most Asian countries, where rice is almost entirely the product of smallholder agriculture, harvesting is carried out manually, although there is a growing interest in mechanical harvesting. Harvesting can be carried out by the farmers themselves, but is also frequently done by seasonal labour groups. Harvesting is followed by threshing, either immediately or within a day or two. Again, much threshing is still carried out by hand but there is an increasing use of mechanical threshers. Subsequently, paddy needs to be dried to bring down the moisture content to no more than 20% for milling. A familiar sight in several Asian countries is paddy laid out to dry along roads. However, in most countries the bulk of drying of marketed paddy takes place in mills, with village-level drying being used for paddy to be consumed by farm families. Mills either sun dry or use mechanical driers or both. Drying has to be carried out quickly to avoid the formation of moulds. Mills range from simple hullers, with a throughput of a couple of tonnes a day, that simply remove the outer husk, to enormous operations that can process 4,000 tonnes a day and produce highly polished rice. A good mill can achieve a paddy-to-rice conversion rate of up to 72% but smaller, inefficient mills often struggle to achieve 60%. These smaller mills often do not buy paddy and sell rice but only service farmers who want to mill their paddy for their own consumption.

Distribution

Because of the importance of rice to human nutrition and food security in Asia, the domestic rice markets tend to be subject to considerable state involvement. While the private sector plays a leading role in most countries, agencies such as BULOG in Indonesia, the NFA in the Philippines, VINAFOOD in Vietnam and the Food Corporation of India are all heavily involved in purchasing of paddy from farmers or rice from mills and in distributing rice to poorer people. BULOG and NFA monopolise rice imports into their countries while VINAFOOD controls all exports from Vietnam.^[67]

Trade

World trade figures are very different to those for production, as only about 5–6% of rice produced is traded internationally. In economic terms, the global rice trade was a small fraction of 1% of world mercantile trade. Many countries consider rice as a strategic food staple, and various governments subject its trade to a wide range of controls and interventions.

Developing countries are the main players in the world rice trade, accounting for 83% of exports and 85% of imports. While there are numerous importers of rice, the exporters of rice are limited. Just five countries – Thailand, Vietnam, China, the United States and India – in decreasing order of exported quantities, accounted for about three-quarters of world rice exports in 2002.^[63] In 2010, the three largest exporters of rice, in decreasing order of quantity exported were Thailand, Vietnam and India. Together, they accounted for nearly 70% of the world rice exports. The primary variety exported by Thailand and Vietnam were Jasmine rice, while exports from India included aromatic Basmati variety. China, an exporter of rice in early 2000s, was a net importer of rice in 2010.^[68] According to a USDA report, the world's largest exporters of rice in 2012 were India (9.75 million tonnes), Vietnam (7 million tonnes), Thailand (6.5 million tonnes), Pakistan (3.75 million tonnes) and the United States (3.5 million tonnes).^[69]

Major importers usually include Nigeria, Indonesia, Bangladesh, Saudi Arabia, Iran, Iraq, Malaysia, the Philippines, Brazil and some African and Persian Gulf countries. Although China and India are the two largest producers of rice in the world, both countries consume the majority of the rice produced domestically, leaving little to be traded internationally.

World's most productive rice farms and farmers

The average world yield for rice was 4.3 tonnes per hectare, in 2010.

Australian rice farms were the most productive in 2010, with a nationwide average of 10.8 tonnes per hectare.^[70]






















Yuan Longping of China National Hybrid Rice Research and Development Center, China, set a world record for rice yield in 2010 at 19 tonnes per hectare on a demonstration plot. In 2011, this record was surpassed by an Indian farmer, Sumant Kumar, with 22.4 tonnes per hectare in Bihar. Both these farmers claim to have employed newly developed rice breeds and System of Rice Intensification (SRI), a recent innovation in rice farming. SRI is claimed to have set new national records in rice yields, within the last 10 years, in many countries. The claimed Chinese and Indian yields have yet to be demonstrated on seven-hectare lots and to be reproducible over two consecutive years on the same farm.^{[71][72][73][74]}

Price

In late 2007 to May 2008, the price of grains rose greatly due to droughts in major producing countries (particularly Australia), increased use of grains for animal feed and US subsidies for bio-fuel production. Although there was no shortage of rice on world markets this general upward trend in grain prices led to panic buying by consumers, government rice export bans (in particular, by Vietnam and India) and inflated import orders by the Philippines marketing board, the National Food Authority. This caused significant rises in rice prices. In late April 2008, prices hit 24 US cents a pound, twice the price of seven months earlier.^[75]

On April 30, 2008, Thailand announced plans for the creation of the Organisation of Rice Exporting Countries (OREC) with the intention that this should develop into a price-fixing cartel for rice.^{[76][77]} However, little progress had been made by mid-2011 to achieve this.

Worldwide consumption

Food consumption of rice by country – 2009 (million metric ton of paddy equivalent) ^[78]	
World Total	531.6
 People's Republic of China	156.3
 India	123.5
 Indonesia	45.3
 Bangladesh	38.2
 Viet Nam	18.4
 Philippines	17.0
 Thailand	13.7
 Japan	10.2
 Burma	10.0
 Brazil	10.0
 Korea	5.8
 Nigeria	4.8
 Egypt	4.6
 Pakistan	4.3
 USA	3.9
 North Korea	3.9
 Nepal	3.5
 Cambodia	3.4
 Sri Lanka	3.2
 Madagascar	3.2
 Malaysia	3.1

As of 2009 world food consumption of rice was 531,639 thousands metric tons of paddy equivalent (354,603 of milled equivalent), while the far largest consumers were China consuming 156,312 thousands metric tons of paddy equivalent (29.4% of the world consumption) and India consuming 123,508 thousands metric tons of paddy equivalent (23.3% of the world consumption).^[78] Between 1961 and 2002, per capita consumption of rice increased by 40%.

Rice is the most important crop in Asia. In Cambodia, for example, 90% of the total agricultural area is used for rice production.^[79]

U.S. rice consumption has risen sharply over the past 25 years, fueled in part by commercial applications such as beer production.^[80] Almost one in five adult Americans now report eating at least half a serving of white or brown rice per day.^[81]

Environmental impacts

Rice cultivation on wetland rice fields is thought to be responsible for 6–29% of the anthropogenic methane emissions annually.^[82] Rice requires slightly more water to produce than other grains.^[83]

Long-term flooding of rice fields cuts the soil off from atmospheric oxygen and causes anaerobic fermentation of organic matter in the soil.^[82] Current contributions of methane from agriculture is ~15% of anthropogenic greenhouse gases, as estimated by the IPCC. Methane is twenty times more potent a greenhouse gas than carbon dioxide.^[84]

A 2010 study found that, as a result of rising temperatures and decreasing solar radiation during the later years of the 20th century, the rice yield growth rate has decreased in many parts of Asia, compared to what would have been observed had the temperature and solar radiation trends not occurred.^{[85][86]} The yield growth rate had fallen 10–20% at some locations. The study was based on records from 227 farms in Thailand, Vietnam, Nepal, India, China, Bangladesh, and Pakistan. The mechanism of this falling yield was not clear, but might involve increased respiration during warm nights, which expends energy without being able to photosynthesize.

Pests and diseases

Rice pests are any organisms or microbes with the potential to reduce the yield or value of the rice crop (or of rice seeds).^[87] Rice pests include weeds, pathogens, insects, nematode, rodents, and birds. A variety of factors can contribute to pest outbreaks, including the overuse of pesticides, improper irrigation, and high rates of nitrogen fertilizer application.^[88] Weather conditions also contribute to pest outbreaks. For example, rice gall midge and army worm outbreaks tend to follow periods of high rainfall early in the wet season, while thrips outbreaks are associated with drought.^[89]

Several nematode species infect rice crops, causing diseases such as Ufra (*Ditylenchus dipsaci*), White tip disease (*Aphelenchoide bessei*), and root knot disease (*Meloidogyne graminicola*). Some nematode species such as *Pratylenchus* spp. are most dangerous in upland rice of all parts of the world. Rice root nematode (*Hirschmanniella* spp.) is a migratory endoparasite which on higher inoculum levels will lead to complete destruction of a rice crop. Beyond being obligate parasites, they also decrease the vigor of plants and increase the plants' susceptibility to other pests and diseases.

Crop protection scientists are trying to develop rice pest management techniques which are sustainable. In other words, to manage crop pests in such a manner that future crop production is not threatened.^[90] Sustainable pest management is based on four principles: biodiversity, host plant resistance (HPR), landscape ecology, and hierarchies in a landscape – from biological to social.^[91] At present, rice pest management includes cultural techniques, pest-resistant rice varieties, and pesticides (which include insecticide). Increasingly, there is evidence that farmers' pesticide applications are often unnecessary, and even facilitate pest outbreaks.^{[92][93][94][95]} By reducing the populations of natural enemies of rice pests,^[96] misuse of insecticides can actually lead to pest outbreaks.^[97] The International Rice Research Institute (IRRI) demonstrated in 1993 that a 87.5% reduction in pesticide use can lead to an overall drop in pest numbers.^[98] IRRI also conducted two campaigns in 1994 and 2003, respectively, which discouraged insecticide misuse and smarter pest management in Vietnam.^{[99][100]}

Rice plants produce their own chemical defenses to protect themselves from pest attacks. Some synthetic chemicals, such as the herbicide 2,4-D, cause the plant to increase the production of certain defensive chemicals and thereby increase the plant's resistance to some types of pests.^[101] Conversely, other chemicals, such as the insecticide imidacloprid, can induce changes in the gene expression of the rice that cause the plant to become more susceptible to attacks by certain types of pests.^[102]

5-Alkylresorcinols are chemicals that can also be found in rice.^[103]


Botanicals, so-called "natural pesticides", are used by some farmers in an attempt to control rice pests. Botanicals include extracts of leaves, or a mulch of the leaves themselves. Some upland rice farmers in Cambodia spread

chopped leaves of the bitter bush (*Chromolaena odorata*) over the surface of fields after planting. This practice probably helps the soil retain moisture and thereby facilitates seed germination. Farmers also claim the leaves are a natural fertilizer and helps suppress weed and insect infestations.^[104]

Among rice cultivars, there are differences in the responses to, and recovery from, pest damage.^{[105][106]} Therefore, particular cultivars are recommended for areas prone to certain pest problems. The genetically based ability of a rice variety to withstand pest attacks is called resistance. Three main types of plant resistance to pests are recognized as nonpreference, antibiosis, and tolerance.^[107] Nonpreference (or antixenosis) describes host plants which insects prefer to avoid; antibiosis is where insect survival is reduced after the ingestion of host tissue; and tolerance is the capacity of a plant to produce high yield or retain high quality despite insect infestation.^[108] Over time, the use of pest resistant rice varieties selects for pests that are able to overcome these mechanisms of resistance. When a rice variety is no longer able to resist pest infestations, resistance is said to have broken down. Rice varieties that can be widely grown for many years in the presence of pests and retain their ability to withstand the pests are said to have durable resistance. Mutants of popular rice varieties are regularly screened by plant breeders to discover new sources of durable resistance.^{[107][109]}

Major rice pests include the brown planthopper^[110] the rice gall midge,^[111] the rice bug,^[105] the rice leafroller, rice weevils, stemborer,^[112] panicle rice mite, rats,^[113] and the weed *Echinochloa crusgali*.^[114]

Major rice diseases include Rice ragged stunt, Sheath Blight, and tungro.^[115] Rice blast, caused by the fungus *Magnaporthe grisea*, is the most significant disease affecting rice cultivation. There is also an ascomycete fungus, *Cochliobolus miyabeanus*, that causes brown spot disease in rice.^{[116][117]}



Chloroxylon is used for Pest Management in Organic Rice Cultivation in Chhattisgarh, India

Parasitic weeds

Rice is parasitized by the weed eudicot *Striga hermonthica*.^[118] This parasitic weed is a devastating pest on the crop. In Rwanda, the national agriculture research institute (ISAR) seeks on a new variety of rice that can resist more to pests and diseases in collaboration with national university of Rwanda at ruhande.

Cultivars

While most rice is bred for crop quality and productivity, there are varieties selected for characteristics such as texture, smell, and firmness. Cultivars exist that are adapted to deep flooding, and these are generally called "floating rice".^[119]

There are four major categories of rice worldwide: indica, japonica, aromatic and glutinous. The different varieties of rice are not considered interchangeable, either in food preparation or agriculture, so as a result, each major variety is a completely separate market from other varieties. It is common for one variety of rice to rise in price while another one drops in price.^[120]

The largest collection of rice cultivars is at the International Rice Research Institute^[121] in the Philippines, with over 100,000 rice accessions^[122] held in the International Rice Genebank.^[123] Rice cultivars are often classified by their grain shapes and texture. For example, Thai Jasmine rice is long-grain and relatively less sticky, as some long-grain rice contains less amylopectin than short-grain cultivars. Chinese restaurants often serve long-grain as plain unseasoned steamed rice though short-grain rice is common as well.

Japanese mochi rice and Chinese sticky rice are short-grain. Chinese people use sticky rice which is properly known as "glutinous rice" (note: glutinous refer to the glue-like characteristic of rice; does not refer to "gluten") to make zongzi. The Japanese table rice is a sticky, short-grain rice. Japanese sake rice is another kind as well.


Indian rice cultivars include long-grained and aromatic Basmati (बासमती) (grown in the North), long and medium-grained Patna rice, and in South India (Andhra Pradesh and Karnataka) short-grained Sona Masuri (also called as Bangaru theegalu). In the state of Tamil Nadu, the most prized cultivar is *ponni* which is primarily grown in the delta regions of the Kaveri River. Kaveri is also referred to as ponni in the South and the name reflects the geographic region where it is grown. In the Western Indian state of Maharashtra, a short grain variety called Ambemohar is very popular. This rice has a characteristic fragrance of Mango blossom.

Aromatic rices have definite aromas and flavors; the most noted cultivars are Thai fragrant rice, Basmati, Patna rice, Vietnamese fragrant rice, and a hybrid cultivar from America, sold under the trade name Texmati. Both Basmati and Texmati have a mild popcorn-like aroma and flavor. In Indonesia, there are also *red* and *black* cultivars.

High-yield cultivars of rice suitable for cultivation in Africa and other dry ecosystems, called the new rice for Africa (NERICA) cultivars, have been developed. It is hoped that their cultivation will improve food security in West Africa.



Rice seed collection from IRRI



Unpolished long-grain rice grains with bran

Draft genomes for the two most common rice cultivars, *indica* and *japonica*, were published in April 2002. Rice was chosen as a model organism for the biology of grasses because of its relatively small genome (~430 megabase pairs). Rice was the first crop with a complete genome sequence.^[124]

On December 16, 2002, the UN General Assembly declared the year 2004 the International Year of Rice. The declaration was sponsored by more than 40 countries.



Polished Indian sona masuri rice grains

Biotechnology

High-yielding varieties

The high-yielding varieties are a group of crops created intentionally during the Green Revolution to increase global food production. This project enabled labor markets in Asia to shift away from agriculture, and into industrial sectors. The first "Rice Car", IR8 was produced in 1966 at the International Rice Research Institute which is based in the Philippines at the University of the Philippines' Los Baños site. IR8 was created through a cross between an Indonesian variety named "Peta" and a Chinese variety named "Dee Geo Woo Gen."^[125]

Scientists have identified and cloned many genes involved in the gibberellin signaling pathway, including GAI1 (Gibberellin Insensitive) and SLR1 (Slender Rice).^[126] Disruption of gibberellin signaling can lead to significantly reduced stem growth leading to a dwarf phenotype. Photosynthetic investment in the stem is reduced dramatically as the shorter plants are inherently more stable mechanically. Assimilates become redirected to grain production, amplifying in particular the effect of chemical fertilizers on commercial yield. In the presence of nitrogen fertilizers, and intensive crop management, these varieties increase their yield two to three times.

Future potential

As the UN Millennium Development project seeks to spread global economic development to Africa, the "Green Revolution" is cited as the model for economic development. With the intent of replicating the successful Asian boom in agronomic productivity, groups like the Earth Institute are doing research on African agricultural systems, hoping to increase productivity. An important way this can happen is the production of "New Rices for Africa" (NERICA). These rices, selected to tolerate the low input and harsh growing conditions of African agriculture, are produced by the African Rice Center, and billed as technology "from Africa, for Africa". The NERICA have appeared in *The New York Times* (October 10, 2007) and *International Herald Tribune* (October 9, 2007), trumpeted as miracle crops that will dramatically increase rice yield in Africa and enable an economic resurgence. Ongoing research in China to develop perennial rice could result in enhanced sustainability and food security.

Golden rice

Rice kernels do not contain vitamin A, so people who obtain most of their calories from rice are at risk of vitamin A deficiency. German and Swiss researchers have genetically engineered rice to produce beta-carotene, the precursor to vitamin A, in the rice kernel. The beta-carotene turns the processed (white) rice a "gold" color, hence the name "golden rice." The beta-carotene is converted to vitamin A in humans who consume the rice.^[127] Although some rice strains produce beta-carotene in the hull, no non-genetically engineered strains have been found that produce beta-carotene in the kernel, despite the testing of thousands of strains. Additional efforts are being made to improve the quantity and quality of other nutrients in golden rice.^[128]

The International Rice Research Institute is currently further developing and evaluating Golden Rice as a potential new way to help address vitamin A deficiency.^[129]

Expression of human proteins

Ventria Bioscience has genetically modified rice to express lactoferrin, lysozyme, and human serum albumin, which are proteins usually found in breast milk. These proteins have antiviral, antibacterial, and antifungal effects.^[130]

Rice containing these added proteins can be used as a component in oral rehydration solutions which are used to treat diarrheal diseases, thereby shortening their duration and reducing recurrence. Such supplements may also help reverse anemia.^[131]

References

- [1] "ProdSTAT" (<http://faostat.fao.org/site/567/DesktopDefault.aspx>). *FAOSTAT*. Retrieved December 26, 2006.
- [2] Smith, Bruce D. *The Emergence of Agriculture*. Scientific American Library, A Division of HPHLP, New York, 1998 ISBN 0-7167-6030-4.
- [3] Why do people throw rice or birdseed as a newly married couple leaves the church? (<http://ask.yahoo.com/20030626.html>). Ask.yahoo.com (June 26, 2003). Retrieved on 2012-05-13.
- [4] International Rice Research Institute *The Rice Plant and How it Grows* (http://www.knowledgebank.irri.org/riceIPM/IPM_Information/PestEcologyBasics/CropGrowthAndPestDamage/RicePlantHowItGrows/The_Rice_plant_and_How_it_Grows.htm) Retrieved January 29, 2008
- [5] rice (<http://www.etymonline.com/index.php?term=rice>), Online Etymology Dictionary
- [6] rice (http://oxforddictionaries.com/definition/rice?rskey=2B5J5N&result=1#m_en_gb0709840), Oxford Dictionaries
- [7] ῥυζα (<http://www.perseus.tufts.edu/hopper/text?doc=Perseus:text:1999.04.0057:entry=o/ruza>), Henry George Liddell, Robert Scott, *A Greek-English Lexicon*, on Perseus
- [8] Witzel, Michael (1999). "Substrate Languages in Old Indo-Aryan" (<http://www.ejvs.laurasianacademy.com/ejvs0501/ejvs0501article.pdf>). *Electronic Journal of Vedic Studies* **5** (1): 26. .
- [9] Thorley, John (1969). "The development of trade between the Roman Empire and the East under Augustus". *Greece & Rome* **16** (2): 222. doi:10.1017/S001738350001706X. JSTOR 642851.
- [10] (2003) *apud* Witzel 2009 (http://www.ias.ac.in/jbiosci/dec2009/Witzel_fulltext), p. 25
- [11] Mayrhofer, Manfred (1996) (in German). *Etymologisches Wörterbuch des Altindoarischen*. **2**. Heidelberg: Universitätsverlag Winter. pp. 598. ISBN 3-8253-4550-5.
- [12] Risks of Talcum Powder (<http://www.preventcancer.com/consumers/cosmetics/talc.htm>). Preventcancer.com. Retrieved on 2012-05-13.
- [13] <http://www.nal.usda.gov/fnic/foodcomp/search/>
- [14] Jianguo G. Wu, Chunhai Shia and Xiaoming Zhanga (2003). "Estimating the amino acid composition in milled rice by near-infrared reflectance spectroscopy". *Field Crops Research* **75**: 1. doi:10.1016/S0378-4290(02)00006-0.
- [15] Kasma Loha-unchit. "White Sticky Rice – Kao Niow" (<http://www.thaifoodandtravel.com/ingredients/sticky.html>). Retrieved October 12, 2012.
- [16] The latter method of using excess water is not desirable with enriched rice, as much of the enrichment additives are flushed away when the water is discarded.
- [17] Watson, p. 15
- [18] Shoichi Ito and Yukihiko Ishikawa Tottori University, Japan. "(Marketing of Value-Added Rice Products in Japan: Germinated Grown Rice and Rice Bread.)" (<http://www.hatsuga.com/DOMER/english/en/GBRRB.html>). Retrieved February 12, 2004.
- [19] "Rice is Life" (<http://www.fao.org/rice2004/en/f-sheet/factsheet3.pdf>). Food and Agricultural Organization of the United Nations. 2004. .
- [20] Bienvenido O. Juliano (1993). "Rice in human nutrition" (<http://www.fao.org/docrep/T0567E/T0567E00.htm>). Food and Agricultural Organization of the United Nations. .
- [21] "Nutrition Info: white rice cooked versus brown rice cooked" (<http://skipthepie.org/cereal-grains-and-pasta/rice-white-long-grain-regular-cooked/compared-to/rice-brown-long-grain-cooked/>). .
- [22] Sungjoon Jang and Zhimin Xu (2009). "Lipophilic and Hydrophilic Antioxidants and Their Antioxidant Activities in Purple Rice Bran" (<http://pubs.acs.org/doi/abs/10.1021/jf803113c>). *Journal of Agricultural and Food Chemistry* **57** (3): 858–862. doi:10.1021/jf803113c. PMID 19138081. .
- [23] Ling, WH; Cheng, QX; Ma, J; Wang, T (2001). "Red and Black Rice Decrease Atherosclerotic Plaque Formation and Increase Antioxidant Status in Rabbits" (<http://jn.nutrition.org/content/131/5/1421.short>). *Journal of Nutrition* **131** (5): 1421–1426. PMID 11340093. .
- [24] "Nutrient data laboratory" (<http://www.nal.usda.gov/fnic/foodcomp/Data/SR18/sr18.html>). United States Department of Agriculture. .
- [25] EFSA Panel on Contaminants in the Food Chain (CONTAM) (October 28, 2009). "Scientific Opinion on Arsenic in Food" (<http://www.efsa.europa.eu/en/efsajournal/pub/1351.htm>). *EFSA Journal* (European Food Safety Authority) **7** (10): 1351. doi:10.2903/j.efsa.2009.1351. . Retrieved November 22, 2012.

- [26] "Arsenic in your food: Our findings show a real need for federal standards for this toxin" (<http://www.consumerreports.org/cro/arsenicinfood.htm>). *Consumer Reports*. November 2012. . Retrieved November 22, 2012.
- [27] FDA.gov - Questions & Answers: FDA's Analysis of Arsenic in Rice and Rice Products (<http://www.fda.gov/Food/FoodSafety/FoodContaminantsAdulteration/Metals/ucm319948.htm>)
- [28] <http://medicalxpress.com/news/2011-12-rice-source-arsenic-exposure.html>
- [29] Consumer Reports Magazine November 2012 - Arsenic in your Food (<http://www.consumerreports.org/cro/magazine/2012/11/arsenic-in-your-food/index.htm#chart>)
- [30] Carol Potera (June 2007). "Food Safety: U.S. Rice Serves Up Arsenic" (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1892142>). *Environmental Health Perspectives* **115** (6): A296. .
- [31] "NHS: Can reheating rice cause food poisoning?" (<http://www.nhs.uk/chq/Pages/can-reheating-rice-cause-food-poisoning.aspx?CategoryID=51&SubCategoryID=215>). .
- [32] IRRI rice knowledge bank (<http://www.knowledgebank.irri.org/>)
- [33] More rice with less water (<http://ciifad.cornell.edu/sri/extmats/philmanual.pdf>) *cornell.edu*. Retrieved on 2012-05-13.
- [34] Plants capable of surviving flooding (<http://www.uu.nl/EN/Current/Pages/Gewaarschuwdeplantenhoudenademinbijoverstroming.aspx>). Uu.nl. Retrieved on 2012-05-13.
- [35] drought tolerance in upland rice (<http://stresstolerant.blogspot.com/2008/09/upland-rice-useful-genetic-resource-for.html>). Stresstolerant.blogspot.com (September 6, 2008). Retrieved on 2012-05-13.
- [36] <http://www.sciencenewline.com/biology/2011050313000047.html>
- [37] Huang, Xuehui; Kurata, Nori; Wei, Xinghua; Wang, Zi-Xuan; Wang, Ahong; Zhao, Qiang; Zhao, Yan; Liu, Kunyan et al. (2012). "A map of rice genome variation reveals the origin of cultivated rice". *Nature*. doi:10.1038/nature11532.
- [38] Vaughan; Lu, B; Tomooka, N (2008). "The evolving story of rice evolution". *Plant Science* **174** (4): 394–408. doi:10.1016/j.plantsci.2008.01.016.
- [39] Harris, David R. (1996). *The Origins and Spread of Agriculture and Pastoralism in Eurasia*. Psychology Press. p. 565. ISBN 1-85728-538-7.
- [40] MacNeish R. S. and Libby J. eds. (1995) *Origins of Rice Agriculture*. Publications in Anthropology No. 13.
- [41] "World's 'oldest' rice found", Dr David Whitehouse" (<http://news.bbc.co.uk/1/hi/sci/tech/3207552.stm>). *BBC News*. October 21, 2003. . Retrieved April 30, 2011.
- [42] Kim, Minkoo (2008). Habu, Junko; Fawcett, Clare; Matsunaga, John M.. ed. *Evaluating multiple narratives: Beyond nationalist, colonialist, imperialist archaeologies*. New York: Springer. p. 128. ISBN 978-0-387-76459-7. "Most scholars were highly skeptical of Lee's report [...] Most specialists agree that rice is not indigenous to the Korean peninsula. The conventional perspective in East Asian archaeology is that rice cultivation started along the banks of the Yangtze River in southern China and subsequently moved northward."
- [43] Kim, Minkoo (2008). "Multivocality, Multifaceted Voices, and Korean Archaeology". *Evaluating Multiple Narratives: Beyond Nationalist, Colonialist, Imperialist Archaeologies*. New York: Springer. p. 118. ISBN 978-0-387-76459-7.
- [44] "Rice's Origins Point to China, Genome Researchers Conclude" (<http://www.sciencenewline.com/biology/2011050313000047.html>). ScienceNewline. . Retrieved December 13, 2011.
- [45] Molina, J.; Sikora, M.; Garud, N.; Flowers, J.M.; Rubinstein, S.; Reynolds, A.; Huang, P.; Jackson, S.; Schaal, B.A.; Bustamante, C.D.; Boyko, A.R.; Purugganan, M.D. (2011). "Molecular evidence for a single evolutionary origin of domesticated rice". *Proceedings of the National Academy of Sciences* **108** (20): 8351–8356. doi:10.1073/pnas.1104686108.
- [46] Maddox, Gregory [ed.] (2006). *Sub-Saharan Africa: An Environmental History*. ABC-CLIO. p. 267. ISBN 1851095551
- [47] National Research Council (1996). "African Rice" (http://books.nap.edu/openbook.php?record_id=2305&page=17). *Lost Crops of Africa: Volume I: Grains* (http://books.nap.edu/openbook.php?record_id=2305). Lost Crops of Africa. **1**. National Academies Press. ISBN 978-0-309-04990-0. . Retrieved July 18, 2008.
- [48] Watson, pp. 17–18
- [49] Robert Sallare, *The Ecology of the Ancient Greek World*, Cornell Univ. Press, 1993. p. 23.
- [50] H.C. Darby (1957). "The face of Europe on the eve of the great discoveries". *The New Cambridge Modern History*. **1**. p. 32. doi:10.1017/CHOL9780521045414.005.
- [51] Rice and Slavery (http://www.slaveryinamerica.org/history/hs_es_rice.htm). Slavery in America. Retrieved on 2012-05-13.
- [52] Carney, Judith Ann (2001). *Black rice: the African origins of rice cultivation in the Americas*. Cambridge: Harvard University Press. ISBN 0-674-00452-3.
- [53] Carolina Gold Rice Foundation (<http://www.carolinagoldricefoundation.org/>). Carolina Gold Rice Foundation. Retrieved on 2012-05-13.
- [54] Farm Raised Crawfish (<http://www.crawfish.com/farmed-crawfish-vs-wild-crawfish.php>). Crawfish.com. Retrieved on 2012-05-13.
- [55] Ching Lee (2005). "Historic Richvale – the birthplace of California rice" (<http://www.cfbf.com/magazine/MagazineStory.cfm?ID=17&ck=70EFD2EC9B086079795C442636B55FB>). California Farm Bureau Federation. . Retrieved August 10, 2007.
- [56] "California's Rice Growing Region" (http://www.calrice.org/e7b_cas_rice_growing_region.htm). California Rice Commission. . Retrieved August 10, 2007.
- [57] Daniel A. Sumner; Henrich Brunke (2003). "The economic contributions of the California rice industry" (http://www.calrice.org/c3a_economic_impact.htm). California Rice Commission. . Retrieved August 10, 2007.
- [58] "Medium Grain Varieties" (http://www.calrice.org/b6a_medium.htm). California Rice Commission. . Retrieved August 10, 2007.

- [59] States Department of Agriculture (<http://www.usda.gov/wps/portal/usdahome?contentidonly=true&contentid=2006/08/0306.xml>) August 2006, Release No. 0306.06, U.S. RICE STATISTICS
- [60] Wadham, Sir Samuel; Wilson, R. Kent and Wood, Joyce; *Land Utilization in Australia*, Melbourne University Press (1957) p. 246
- [61] Australian Bureau of Meteorology; *Climatic Atlas of Australia: Rainfall*, Bureau of Meteorology, Melbourne, Victoria (2000)
- [62] fao.org (FAOSTAT). "Countries by commodity (Rice, paddy)" (<http://faostat.fao.org/site/339/default.aspx>). . Retrieved May 26, 2012.
- [63] "Sustainable rice production for food security" (<http://www.fao.org/DOCREP/006/Y4751E/y4751e00.htm>). Food and Agriculture Organization of the United Nations. 2003. .
- [64] "World Wheat, Corn and Rice" (http://nue.okstate.edu/crop_information/world_wheat_production.htm). Oklahoma State University, FAOSTAT. .
- [65] "MISSING FOOD: The Case of Postharvest Grain Losses in Sub-Saharan Africa" (http://siteresources.worldbank.org/INTARD/Resources/MissingFoods10_web.pdf). The World Bank. April 2011. .
- [66] Basavaraja, H.; Mahajanashetti, S.B.; Udagatti, Naveen C. "Economic Analysis of Post-harvest Losses in Food Grains in India: A Case Study of Karnataka" (<http://ageconsearch.umn.edu/bitstream/47429/2/8.pdf>). *Agricultural Economics Research Review* **20**: 117–126. .
- [67] Shahidur Rashid, Ashok Gulari and Ralph Cummings Jnr (eds), *From Parastatals to Private Trade*. International Food Policy Research Institute and Johns Hopkins University Press, 2008 ISBN 0-8018-8815-8
- [68] "World Rice Trade" (<http://www.fas.usda.gov/psdonline/psdreport.aspx?hidReportRetrievalName=BVS&hidReportRetrievalID=677&hidReportRetrievalTemplateID=7>). United States Department of Agriculture. November 2011. .
- [69] <http://www.financialexpress.com/news/india-is-worlds-largest-rice-exporter-usda/1023491/0>
- [70] "FAOSTAT: Production-Crops, 2010 data" (<http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567#anchor>). Food and Agriculture Organization of the United Nations. 2011. .
- [71] L.P. Yuan (2010). "A Scientist's Perspective on Experience with SRI in CHINA for Raising the Yields of Super Hybrid Rice" (http://ciifad.cornell.edu/sri/proc1/sri_06.pdf). .
- [72] "Indian farmer sets new world record in rice yield" (<http://www.philstar.com/Article.aspx?articleId=759471>). The Philippine Star. December 18, 2011. .
- [73] "Grassroots heroes lead Bihar's rural revolution" (<http://indiatoday.intoday.in/story/grassroots-heroes-lead-rural-revolution-of-bihar/1/168028.html>). India Today. January 10, 2012. .
- [74] "System of Rice Intensification" (<http://sri.ciifad.cornell.edu/>). Cornell University. 2011. .
- [75] "Cyclone fuels rice price increase" (<http://news.bbc.co.uk/2/hi/business/7387251.stm>). *BBC News*, May 7, 2008
- [76] "Mekong nations to form rice price-fixing cartel" (<http://www.radioaustralia.net.au/news/stories/200804/s2232076.htm?tab=asia>), Radio Australia, April 30, 2008.
- [77] "PM floats idea of five-nation rice cartel" (http://www.bangkokpost.com/News/01May2008_news07.php), *Bangkok Post*, May 1, 2008.
- [78] FAO (FAOSTAT). "Food Balance Sheets > Commodity Balances > Crops Primary Equivalent" (<http://faostat.fao.org/site/616/DesktopDefault.aspx?PageID=616#anchor>). . Retrieved August 17, 2012.
- [79] Don Puckridge, "The Burning of the Rice" (<http://sidharta.com/books/index.jsp?uid=67>). Sidharta.com. Retrieved on 2012-05-13.
- [80] United States Department of Agriculture (USDA) Economic Research Service. "Briefing Rooms: Rice" (<http://www.ers.usda.gov/Briefing/Rice/>). . Retrieved April 24, 2008.
- [81] Iowa State University (July 2005). *Rice Consumption in the United States: New Evidence from Food Consumption Surveys* (<http://publications.iowa.gov/2781/>). .
- [82] Neue Heinz-Ulrich (1993). "Methane emission from rice fields: Wetland rice fields may make a major contribution to global warming" (<http://www.ciesin.org/docs/004-032/004-032.html>). *BioScience* **43** (7): 466–73. .
- [83] report12.pdf Virtual Water Trade – Proceedings of the International Expert Meeting on Virtual Water Trade (<http://www.waterfootprint.org/Reports/Report12.pdf>), p. 108
- [84] IPCC. Climate Change 2007: Synthesis Report (http://www.ipcc.ch/pdf/assessment-report/ar4/syr/ar4_syr.pdf). United Nations Environment Programme, 2007:Ch5, 8, and 10. IPCC.ch
- [85] Welch, Jarrod R.; Vincent, J.R.; Auffhammer, M.; Doberman, A.; Moya, P.; Dawe, D. (2010). "Rice yields in tropical/subtropical Asia exhibit large but opposing sensitivities to minimum and maximum temperatures". *Proc. Natl. Acad. Sci. U.S.A.* **107** (33): 14562–7. doi:10.1073/pnas.1001222107. PMC 2930450. PMID 20696908.
- [86] Black, Richard Rice yields falling under global warming (<http://www.bbc.co.uk/news/science-environment-10918591>) BBC News Science & Environment (August 9, 2010). Retrieved August 9, 2010.
- [87] Gary C. Jahn; JA Litsinger, Y Chen and A Barrion (2007). "Integrated Pest Management of Rice: Ecological Concepts". In O Koul and GW Cuperus. *Ecologically Based Integrated Pest Management*. CAB International. pp. 315–366. ISBN 978-1-84593-064-6.
- [88] Jahn, Gary C.; Almazan, Liberty P.; Pacia, Jocelyn B. (2005). "Effect of Nitrogen Fertilizer on the Intrinsic Rate of Increase of *Hysteronura setariae* (Thomas) (Homoptera: Aphididae) on Rice (*Oryza sativa* L.)" (<http://docserver.esa.catchword.org/deliver/cw/pdf/esa/freepdfs/0046225x/v34n4s26.pdf>). *Environmental Entomology* **34** (4): 938. doi:10.1603/0046-225X-34.4.938. .
- [89] Douangboupha, B, K Khamphoukeo, S Inthavong, J Schiller, and GC Jahn. 2006. Pests and diseases of the rice production systems of Laos ([http://aci.gov.au/files/node/756/Rice In Laos chapter 16-25.pdf](http://aci.gov.au/files/node/756/Rice%20In%20Laos%20chapter%2016-25.pdf)). Chapter 17, pp. 265–281. In JM Schiller, MB Chanphengxay, B Linqvist, and S Appa Rao, editors. *Rice in Laos*. Los Baños (Philippines): IRRI. ISBN 978-971-22-0211-7.
- [90] Jahn, GC, B. Khiev, C Pol, N Chhorn, S Pheng, and V Preap. 2001. Developing sustainable pest management for rice in Cambodia. pp. 243–258, In S. Suthipradit, C. Kuntha, S. Lorlowhakarn, and J. Rakngan [eds.]. "Sustainable Agriculture: Possibility and Direction" Bangkok

- (Thailand): National Science and Technology Development Agency.
- [91] Savary, S.; Horgan, F.; Willocquet, L.; Heong, K.L. (2012). "A review of principles for sustainable pest management in rice". *Crop Protection* **32**: 54. doi:10.1016/j.cropro.2011.10.012.
- [92] Jahn, GC, S Pheng, B Khiev, and C Pol. 1996. Farmers' pest management and rice production practices in Cambodian lowland rice. Cambodia-IRRI-Australia Project (CIAP), Baseline Survey Report No. 6. CIAP Phnom Penh, Cambodia.
- [93] Bangladeshi farmers banish insecticides (<http://www.scidev.net/Features/index.cfm?fuseaction=readfeatures&itemid=306&language=1>). SCIDEV.net (July 30, 2004). Retrieved on 2012-05-13.
- [94] IRRI.org (<http://www.youtube.com/watch?v=VKB-pg8EGMQ>). Youtube.com (November 20, 2006). Retrieved on 2012-05-13.
- [95] Wang, Li-Ping; Shen, Jun; Ge, Lin-Quan; Wu, Jin-Cai; Yang, Guo-Qin; Jahn, Gary C. (2010). "Insecticide-induced increase in the protein content of male accessory glands and its effect on the fecundity of females in the brown planthopper *Nilaparvata lugens* Stål (Hemiptera: Delphacidae)". *Crop Protection* **29** (11): 1280. doi:10.1016/j.cropro.2010.07.009.
- [96] Jahn, G.C. (1992). "Rice pest control and effects on predators in Thailand". *Insecticide & Acaricide Tests* **17**: 252–3.
- [97] Cohen, J. E., Schoenly, K., Heong, K. L., Justo, H., Arida, G., Barrion, A. T., & Litsinger, J. A. (1994). "A Food-Web Approach to Evaluating the Effect of Insecticide Spraying on Insect Pest Population-Dynamics in a Philippine Irrigated Rice Ecosystem" (<http://www.mendeley.com/research/a-foodweb-approach-to-evaluating-the-effect-of-insecticide-spraying-on-insect-pest-population-dynamics-in-a-philippine-irrigated-rice-ecosystem/>). *Journal of Applied Ecology* **31-issue=4**: 747–763. .
- [98] (<http://irri.org/knowledge/publications/rice-today/special-reports/science-shorts/the-pesticide-paradox>)
- [99] 'Three Gains, Three Reductions' (<http://ricehoppers.net/2010/10/three-reduction-three-gains-a-factor-in-vietnam-s-continuing-success-says-thai-report/>). Ricehoppers.net (October 12, 2010). Retrieved on 2012-05-13.
- [100] 'No Early Spray' (<http://ricehoppers.net/wp-content/uploads/2010/04/Escalada-et-al-2009.pdf>) *ricehoppers.net* (April 2010). Retrieved on 2012-05-13.
- [101] Xin, Zhaojun; Yu, Zhaonan; Erb, Matthias; Turlings, Ted C. J.; Wang, Baohui; Qi, Jinfeng; Liu, Shengning; Lou, Yonggen (2012). "The broad-leaf herbicide 2,4-dichlorophenoxyacetic acid turns rice into a living trap for a major insect pest and a parasitic wasp". *New Phytologist* **194** (2): 498–510. doi:10.1111/j.1469-8137.2012.04057.x. PMID 22313362.
- [102] Cheng, Yao; Shi, Zhao-Peng; Jiang, Li-Ben; Ge, Lin-Quan; Wu, Jin-Cai; Jahn, Gary C. (2012). "Possible connection between imidacloprid-induced changes in rice gene transcription profiles and susceptibility to the brown plant hopper *Nilaparvata lugens* Stål (Hemiptera: Delphacidae)". *Pesticide Biochemistry and Physiology* **102** (3): 213–219. doi:10.1016/j.pestbp.2012.01.003. PMC 3334832. PMID 22544984.
- [103] Yoshikatsu Suzuki, , Minoru Kurano, Yasuaki Esumi, Isamu Yamaguchi and Yoshiharu Doi (2003). "Biosynthesis of 5-alkylresorcinol in rice: incorporation of a putative fatty acid unit in the 5-alkylresorcinol carbon chain". *Bioorganic Chemistry* **31** (6): 437–452. doi:10.1016/j.bioorg.2003.08.003. PMID 14613765.
- [104] Jahn, GC, C Pol, B Khiev, S Pheng, and N Chhorn. 1999. Farmer's pest management and rice production practices in Cambodian upland and deepwater rice. Cambodia-IRRI-Australia Project, Baseline Survey Rpt No. 7
- [105] Jahn, GC; Domingo, I; Almazan, ML; Pacia, J; Pacia, Jocelyn (2004). "Effect of rice bug *Leptocorisa oratorius* (Hemiptera: Alydidae) on rice yield, grain quality, and seed viability". *Journal of economic entomology* **97** (6): 1923–7. doi:10.1603/0022-0493-97.6.1923. PMID 15666746.
- [106] Khiev, B.; Jahn, G.C.; Pol, C.; Chhorn N. (2000). "Effects of simulated pest damage on rice yields" (<http://www.mendeley.com/research/effects-of-simulated-pest-damage-on-rice-yields/>). *IRRN* **25** (3): 27–8. .
- [107] Sangha, Jatinder Singh; Chen, Yolanda H.; Palchamy, Kadirvel; Jahn, Gary C.; Maheswaran, M.; Adalla, Candida B.; Leung, Hei (2008). "Categories and Inheritance of Resistance to *Nilaparvata lugens* (Hemiptera: Delphacidae) in Mutants of Indica Rice 'IR64'". *Journal of Economic Entomology* **101** (2): 575–83. doi:10.1603/0022-0493(2008)101[575:CAIORT]2.0.CO;2. ISSN 0022-0493. PMID 18459427.
- [108] Kogan, M.; Ortman, E.F. (1978). "Antixenosis a new term proposed to defined to describe Painter's "non-preference" modality of resistance". *Bull. Entomol. Soc. Am.* **24**: 175–6.
- [109] Liu, L., Z. Van, Q. Y. Shu, and M. Maluszynski (2004). "Officially released mutant varieties in China". *Mutat. Breed. Rev* **14** (1): 64.
- [110] V Preap, MP Zalucki, and GC Jahn (2006). "Brown planthopper outbreaks and management" (<http://www.cardi.org.kh/images/stories/CJA/Vol07-N01-Jan-Jun-2006.pdf>). *Cambodian Journal of Agriculture* **7** (1): 17–25. .
- [111] J. Bennett, JS Bentur, IC Pasula, K. Krishnaiah, [eds]. New approaches to gall midge resistance in rice. International Rice Research Institute and Indian Council of Agricultural Research.
- [112] IRRI Rice insect pest factsheets (<http://www.knowledgebank.irri.org/ipm/index.php/factsheets>)
- [113] G Singleton, L Hinds, H Leirs and Zhibin Zhang [Eds.] "Ecologically-based rodent management" ACIAR, Canberra. Ch. 17, pp. 358–371 ISBN 1-86320-262-5.
- [114] Pheng, S, B Khiev, C Pol and GC Jahn (2001). "Response of two rice cultivars to the competition of *Echinochloa crus-gali* (L.) P. Beauv" (<http://ejournals.ph/index.php?journal=IRRN&page=article&op=view&path=[]=4318>). *International Rice Research Institute Notes (IRRN)* **26** (2): 36–37. .
- [115] IRRI Rice Diseases factsheets (<http://www.knowledgebank.irri.org/ipm/index.php/diseases-crop-health-2733>). Knowledgebank.irri.org. Retrieved on 2012-05-13.
- [116] Rice Brown Spot: essential data (<http://www.cbwinfo.com/Biological/PlantPath/CM.html>). CBWinfo.com. Retrieved on 2012-05-13.

- [117] *Cochliobolus* (<http://www.invasive.org/browse/tax.cfm?fam=683&genus=Cochliobolus>). Invasive.org (May 4, 2010). Retrieved on 2012-05-13.
- [118] Yoshida, Satoko; Maruyama, Shinichiro; Nozaki, Hisayoshi (2010). "Horizontal Gene Transfer by the Parasitic Plant *Stiga hermanthica*". *Science* **328** (5982): 1128. doi:10.1126/science.1187145. PMID 20508124.
- [119] Rice (<http://www.cgiar.org/impact/research/rice.html>). Cgiar.org. Retrieved on 2012-05-13.
- [120] "The U.S. Rice Export Market" (<http://www.ers.usda.gov/briefing/Rice/SpecialArticle/USricemarket.pdf>). USDA. November 2000. .
- [121] "Home" (<http://www.irri.org>). Irri.org. . Retrieved 2012-08-13.
- [122] IRRRI.org (<http://www.irri.org/media/press/press.asp?id=1>)
- [123] Jackson MT (September 1997). "Conservation of rice genetic resources: the role of the International Rice Genebank at IRRI" (<http://www.kluweronline.com/art.pdf?issn=0167-4412&volume=35&page=61>). *Plant Mol. Biol.* **35** (1–2): 61–7. doi:10.1023/A:1005709332130. PMID 9291960. .
- [124] Gillis, Justing (August 11, 2005). "Rice Genome Fully Mapped" (<http://www.washingtonpost.com/wp-dyn/content/article/2005/08/10/AR2005081001054.html?referrer=email>). *Washington Post*. .
- [125] Rice Varieties: IRRI Knowledge Bank. Accessed August 2006 (<http://www.knowledgebank.irri.org/regionalSites/nepal/RiceVarieties.htm>)
- [126] Yamaguchi, S. (2008). "Gibberellin Metabolism and its Regulation". *Ann Rev Plant Biol* **59**: 225–251. doi:10.1146/annurev.arplant.59.032607.092804.
- [127] "Researchers Determine That Golden Rice Is an Effective Source of Vitamin A" (<http://www.goldenrice.org/PDFs/ASNonGR.pdf>). American Society for Nutrition. 2009. . Retrieved October 28, 2010.
- [128] Grand Challenges in Global Health, Engineering Rice for High Beta Carotene, Vitamin E and Enhanced Fe and Zn Bioavailability (<http://www.grandchallenges.org/ImproveNutrition/Challenges/NutrientRichPlants/Pages/Rice.aspx>), grandchallenges.org. Retrieved April 14, 2012
- [129] International Rice Research Institute: Golden Rice (http://www.irri.org/index.php?option=com_k2&view=itemlist&task=category&id=764:golden-rice-at-irri&lang=en)
- [130] Emma Marris (May 18, 2007). "Rice with human proteins to take root in Kansas". *Nature*. doi:10.1038/news070514-17.
- [131] Bethell DR, Huang J (2004). "Recombinant human lactoferrin treatment for global health issues: iron deficiency and acute diarrhea". *Biometals* **17** (3): 337–42. doi:10.1023/B:BIOM.0000027714.56331.b8. PMID 15222487.

Bibliography

- Watson, Andrew (1983). *Agricultural innovation in the early Islamic world*. Cambridge University Press ISBN 0-521-06883-5.
- Andrew A. Meharg and Fang-Jie Zhao, *Arsenic & Rice*, Springer (February 29, 2012), hardcover, 182 pages, ISBN-10: 9400729464 ISBN-13: 978-9400729469

External links

- Rice (http://www.dmoz.org/Science/Agriculture/Field_Crops/Cereals/Rice/) at the Open Directory Project
- International Rice Research Institute (<http://irri.org/>)
- Rice latest trade data on ITC Trade Map (http://www.trademap.org/open_access/Index.aspx?proceed=true&product=1006)
- Global Rice Science Partnership (<http://irri.org/our-science/global-rice-science-partnership-grisp>)
- Rice Today magazine (<http://irri.org/ricetoday>)
- Calories in rice (<http://thecaloriecounter.net/calories-in-rice.html/>)
- Interviews with Pioneers of Rice Research (http://archive.irri.org/publications/today/Pioneer_Interviews.asp)
- Rice Knowledge Bank (<http://irri.org/knowledge/irri-training/knowledge-bank>)
- A Brief History of Rice (p. 9 – p. 12) (http://www.pub.ac.za/resources/docs/biotech_biodiv_module_gr11_2005.pdf)
- A Cuban Skirmish for Rice (<http://www.havanatimes.org/?p=25078>) by Isbel Diaz Torres, *Havana Times*, June 16, 2010
- National Food Authority. (<http://nfa.gov.ph/>)
- Philippine Rice Research Institute. (<http://philrice.gov.ph/>)
- Safe Storage of Cooked Rice (<http://www.abc.net.au/health/talkinghealth/factbuster/stories/2009/01/27/2475255.htm>)

- Rice Research and Practice (<http://uark.libguides.com/rice>)
-

Zeiden, Zeimusu, Zfr, ZhongHan, Zocky, Zollerrria, Zondor, Zykan, Zzorse, Zzuuzz, Zæren Möck, 5 מרכבה גדול, كاشف عقيل, 大西洋鲑, 리듬, 2399 anonymous edits

Image Sources, Licenses and Contributors

File:Koeh-232.jpg *Source:* <http://en.wikipedia.org/w/index.php?title=File:Koeh-232.jpg> *License:* unknown *Contributors:* -

File:Nihonbashi bridge in Edo.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Nihonbashi_bridge_in_Edo.jpg *License:* Public Domain *Contributors:* Morio, Petrusbarbygere, Red devil 666, Tokorokoko, Zolo, 1 anonymous edits

File:Watermill at Onden.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Watermill_at_Onden.jpg *License:* Public Domain *Contributors:* Anna reg, Petrusbarbygere, Red devil 666, Tetrís L, Tokorokoko, Zolo, 立花左近, 1 anonymous edits

File:Rice plantation in Java.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Rice_plantation_in_Java.jpg *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* Gunkarta Gunawan Kartapranata

File:Rice 02.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Rice_02.jpg *License:* Creative Commons Attribution 2.0 *Contributors:* Fg2, Gryffindor, Mattes, Orrling, Ranveig, Spolloman, TwoWings, Vmenkov, 1 anonymous edits

File:Bbbulacancoolwavesjf.JPG *Source:* <http://en.wikipedia.org/w/index.php?title=File:Bbbulacancoolwavesjf.JPG> *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* User:Ramon FVelasquez

File:Niyamgiri rice.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Niyamgiri_rice.jpg *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* User:Pankaj Oudhia

File:Rice p1160004.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Rice_p1160004.jpg *License:* Creative Commons Attribution-ShareAlike 3.0 Unported *Contributors:* User:David.Monniaux

File:Chinese Chawal in Basmati.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Chinese_Chawal_in_Basmati.jpg *License:* Creative Commons Attribution-ShareAlike 3.0 Unported *Contributors:* Khalid Mahmood

File:Riceandcereal.JPG *Source:* <http://en.wikipedia.org/w/index.php?title=File:Riceandcereal.JPG> *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* Jon Ascton at en.wikipedia

File:Kome.JPG *Source:* <http://en.wikipedia.org/w/index.php?title=File:Kome.JPG> *License:* GNU Free Documentation License *Contributors:* JackyR, Kri, Maksim, Namazu-tron, Orrling, トクト, 1 anonymous edits

File:Japanreis.JPG *Source:* <http://en.wikipedia.org/w/index.php?title=File:Japanreis.JPG> *License:* GNU Free Documentation License *Contributors:* Okapi

File:RiceStems400x1.jpg *Source:* <http://en.wikipedia.org/w/index.php?title=File:RiceStems400x1.jpg> *License:* Creative Commons Attribution-Share Alike *Contributors:* John Alan Elson

File:Rice Animation.gif *Source:* http://en.wikipedia.org/w/index.php?title=File:Rice_Animation.gif *License:* Public Domain *Contributors:* Namazu-tron

File:Medicinal Rice.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Medicinal_Rice.jpg *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* User:Pankaj Oudhia

File:Ancient rice.png *Source:* http://en.wikipedia.org/w/index.php?title=File:Ancient_rice.png *License:* Creative Commons Attribution 2.5 *Contributors:* Ayacop, Daniel Mietchen, Masur, NeverDoING, Snek01, Sreejithk2000, Überraschungsbilder

File:Rice-fields-Indonesia-(Java).jpg *Source:* [http://en.wikipedia.org/w/index.php?title=File:Rice-fields-Indonesia-\(Java\).jpg](http://en.wikipedia.org/w/index.php?title=File:Rice-fields-Indonesia-(Java).jpg) *License:* Creative Commons Attribution-ShareAlike 1.0 Generic *Contributors:* Indon, Jurema Oliveira, KTo288, Vadakkan

File:Paddy West Bengal.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Paddy_West_Bengal.jpg *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* User:Amartyabag

File:Banaue Rice Terrace Close Up (2).JPG *Source:* [http://en.wikipedia.org/w/index.php?title=File:Banaue_Rice_Terrace_Close_Up_\(2\).JPG](http://en.wikipedia.org/w/index.php?title=File:Banaue_Rice_Terrace_Close_Up_(2).JPG) *License:* Public domain *Contributors:* Agricmarketing at en.wikipedia

File:burmarice.JPG *Source:* <http://en.wikipedia.org/w/index.php?title=File:Burmarice.JPG> *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* Ericwinny

File:Ambositra 04.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Ambositra_04.jpg *License:* Creative Commons Attribution-Sharealike 3.0,2.5,2.0,1.0 *Contributors:* Bernard Gagnon

File:MansfieldWinnowingBarn.jpg *Source:* <http://en.wikipedia.org/w/index.php?title=File:MansfieldWinnowingBarn.jpg> *License:* GNU Free Documentation License *Contributors:* Thomas Namey, [www.nameydesign.com Namey Design Studios], aka Namey Design Studios at en.wikipedia

File:Flag of the People's Republic of China.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_the_People's_Republic_of_China.svg *License:* Public Domain *Contributors:* Drawn by User:SKopp, redrawn by User:Denelson83 and User:Zscout370 Recode by cs:User:xfi- (code), User:Shizhao (colors)

File:Flag of India.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_India.svg *License:* Public Domain *Contributors:* Anomie, Mifter

File:Flag of Indonesia.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Indonesia.svg *License:* Public Domain *Contributors:* Drawn by User:SKopp, rewritten by User:Gabbe

File:Flag of Bangladesh.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Bangladesh.svg *License:* Public Domain *Contributors:* User:SKOpp

File:Flag of Vietnam.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Vietnam.svg *License:* Public Domain *Contributors:* Lưu Ly về lại theo nguồn trên

File:Flag of Myanmar.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Myanmar.svg *License:* Public Domain *Contributors:* *drew, AnonMoos, CommonsDelinker, Duduziq, Fry1989, Gunkarta, Homo lupus, Idh0854, Josegeographic, Klemen Kocjancic, Legnaw, Mason Decker, Mattes, Neq00, Nightstallion, Pixeltoo, Rfc1394, Rodejong, SeNeKa, Stevanb, ThomasPusch, UnreifeKirsche, VIVIDUPPERS, WikipediaMaster, Winzipas, Xiengyod, Zscout370, 白布飘扬, 10 anonymous edits

File:Flag of Thailand.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Thailand.svg *License:* Public Domain *Contributors:* Zscout370

File:Flag of the Philippines.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_the_Philippines.svg *License:* Public Domain *Contributors:* User:Achim1999

File:Flag of Brazil.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Brazil.svg *License:* Public Domain *Contributors:* Anomie

File:Flag of the United States.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_the_United_States.svg *License:* Public Domain *Contributors:* Anomie

File:Flag of Japan.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Japan.svg *License:* Public Domain *Contributors:* Anomie

File:Flag of Cambodia.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Cambodia.svg *License:* Public Domain *Contributors:* Open Clip Art Library, first uploaded by Nightstallion; redraw the towers of Angkor Wat by User:Xiengyod.

File:Flag of Pakistan.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Pakistan.svg *License:* Public Domain *Contributors:* User:Zscout370

File:Flag of South Korea.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_South_Korea.svg *License:* Public Domain *Contributors:* Various

File:Flag of Madagascar.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Madagascar.svg *License:* Public Domain *Contributors:* User:SKOpp

File:Flag of Egypt.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Egypt.svg *License:* Public Domain *Contributors:* Open Clip Art

File:Flag of Sri Lanka.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Sri_Lanka.svg *License:* Public Domain *Contributors:* Zscout370

File:Flag of Nepal.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Nepal.svg *License:* Public Domain *Contributors:* Drawn by User:Pumbaa80, User:Achim1999

File:Flag of Nigeria.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Nigeria.svg *License:* Public Domain *Contributors:* User:Jhs

File:Flag of Laos.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Laos.svg *License:* Public Domain *Contributors:* User:SKOpp

Image:RiceYield.png *Source:* <http://en.wikipedia.org/w/index.php?title=File:RiceYield.png> *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* AndrewMT

File:Flag of Korea 1882.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Korea_1882.svg *License:* GNU General Public License *Contributors:* Daniil Ivanov

File:Flag of North Korea.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_North_Korea.svg *License:* Public Domain *Contributors:* Zscout370

File:Flag of Malaysia.svg *Source:* http://en.wikipedia.org/w/index.php?title=File:Flag_of_Malaysia.svg *License:* Public Domain *Contributors:* Achim1999, Ah Cong Strike, AnonMoos, Arteyu, Avala, Cyn, DarknessVisitor, Duduziq, Er Komandante, Fibonacci, Fred J, Fry1989, Herbythyme, Homo lupus, Juiced lemon, Klemen Kocjancic, Ludger1961, Morio, Nick, Reisio, Rocket000, SKOpp, Sarang, Tryphon, VAO HK, Zscout370, 白布飘扬, 20 anonymous edits

File:Chloroxylon from India.JPG *Source:* http://en.wikipedia.org/w/index.php?title=File:Chloroxylon_from_India.JPG *License:* Creative Commons Attribution-Sharealike 3.0 *Contributors:* User:Pankaj Oudhia

File:Rice diversity.jpg *Source:* http://en.wikipedia.org/w/index.php?title=File:Rice_diversity.jpg *License:* Creative Commons Attribution 2.0 *Contributors:* IRRI Images
File:Unpolished-rice.jpg *Source:* <http://en.wikipedia.org/w/index.php?title=File:Unpolished-rice.jpg> *License:* GNU Free Documentation License *Contributors:* Sanjay Acharya
File:sona-masuri.jpg *Source:* <http://en.wikipedia.org/w/index.php?title=File:Sona-masuri.jpg> *License:* GNU Free Documentation License *Contributors:* Sanjay Acharya (Sanjay ach at en.wikipedia)

License

Creative Commons Attribution-Share Alike 3.0 Unported
[//creativecommons.org/licenses/by-sa/3.0/](https://creativecommons.org/licenses/by-sa/3.0/)
