




Greener Journal of Agricultural Sciences

ISSN: 2276-7770 Impact Factor 2012 (UJRD): 0.7904 ICV 2012: 6.15

Potency of Some Plant Extracts and Pesticides on Bacterial Leaf Blight Diseases of Cocoyam (*Colocasia Esculenta*) in Umudike, South Eastern Nigeria

By

**Emma Opara
Theresa C. Njoku
Chikodi Isaiah**



Research Article

Potency of Some Plant Extracts and Pesticides on Bacterial Leaf Blight Diseases of Cocoyam (*Colocasia Esculenta*) in Umudike, South Eastern Nigeria

Emma Opara*¹, Theresa C. Njoku² and Chikodi Isaiah³

^{1,3}Department of Plant Health Management, Michael Okpara University of Agriculture, Umudike.

²Department of Crop Science, Imo State University, Owerri.

¹*Corresponding Author's Email: euopara22@yahoo.com, Tel: 08035761578.

ABSTRACT

A pot trial was conducted in 2010 and 2011 cropping seasons at the screen house of Michael Okpara University of Agriculture, Umudike to determine the potency of some plant extracts and pesticides for the control of bacterial leaf blight of cocoyam (*Colocasia esculenta* L.) Scholt. The materials employed include: lemon grass (*Cymbopogon citrates*), Black pepper seed (*Piper guineense*), orange peel (*Citrus sinensis*), streptomycin sulphate (antibiotic), copper oxychloride (fungicide) and sterile water (control). The experiment was laid in complete randomized design (CRD) and replicated six times. The plant materials used were first made into aqueous extracts using cold water extraction and diluted to 10% concentration. Application on leaves was by use of a hand sprayer for two weekly intervals. Data were collected on disease severity and incidence including growth parameters and on yield of corms. The results obtained showed that all the plant extracts were able to reduce disease and increase yield better than the untreated control.

Keywords: *Colocasia esculenta*, bacterial leaf blight, plant extracts, *Xanthomonas campestris* pv. *Campestris*.

INTRODUCTION

Cocoyam *Colocasia esculenta* (L.) Scholt constitutes one of the basic food crops of major economic importance in the south eastern Nigeria. It ranks the third after cassava and yam, in terms of total production, land area under crop and consumption (Chukwu and Nwosu, 2008). There are two main edible types of cocoyam in Nigeria viz: *Colocasia esculenta* (L.) Scholt, otherwise known as 'taro' and *Xanthosoma sagittifolium* also known as 'tannia' (Arene and Ene, 1987). The former is by far more popular than the later. Taro requires heavy fertile upland soil and plentiful rainfall for good yield. It does well also in a fertile low land environment (Agboola, 1987; Anikwe *et al.*, 2007) and commonly grown in South Eastern zone of Nigeria. On the other hand, tannia (*Xanthosoma sagittifolium*) is well cultivated in the South Western zone of Nigeria (Ikwelle, 1999).

Currently, *Colocasia esculenta* (taro) is seriously threatened to extinction in south eastern Nigeria as a result of its high susceptibility to bacterial leaf blight disease. Therefore this study was undertaken to isolate and identify the causal organism and determine the potency of some selected local plant extracts in the control of bacterial leaf blight of *Colocasia esculenta*.

MATERIALS AND METHODS

The study was carried out for two years during the 2010 and 2011 cropping seasons under pot trials. Top soil was collected from the Eastern farm site of the Michael Okpara University of Agric, Umudike. Soil analysis and characterization was conducted to determine the macro and micro elements and other constituents of the soil. The soil was then poured into a cut drum, moistened and the temperature raised until it reached a temperature of 80°C for 20minutes for proper sterilization, and then allowed to cool down before mixing with poultry droppings and fine soil (to improve its porosity) in the ratio of 3: 2: 1 of top soil, fine sand and poultry droppings respectively (the poultry dropping was allowed to decompose for two weeks before use) and later the sterilized soil mixture dispensed into plastic pots of 15cm in diameter (three quarter filled).

The experiment was arranged in a completely randomized design (CRD) consisting of six treatments and six replications. The plant materials used as extracts were lemon grass (*Cymbopogon citrates*) black pepper seed (*Piper guineense*), orange peel (*Citrus sinensis*) and two synthetic pesticides namely streptomycin sulphate and copper oxychloride. Cocoyam corms were planted into experimental pots at the rate of one seedling per pot.

PREPARATION OF PLANT EXTRACTS

The plant materials collected were washed and air dried and 50g weighed out before grinding them into powder for use in this study. The powdered material was suspended in 100ml of sterile water and later diluted to 20% concentration, allowed to stand in water for 24 hours after which the suspension was filtered using cheese cloth and the filtrate obtained used as cold water aqueous extracts diluted (Stoll, 2000; Amadioha, 2003). Also two synthetic pesticides, which included streptomycin sulphate (antibiotic) and copper oxychloride (fungicide) were dissolved in cold sterile water to obtain, 1% and 10% respectively which served as positive checks (Whitehead, 1999), also sterile water was served as control experiment.

PREPARATION OF BACTERIAL INOCULUM

Infected leaf samples were collected from the university farms and brought into the laboratory, thoroughly washed with sterile water and were surface sterilized for 10 seconds in (0.5%) aqueous solution of sodium hypochlorite, the leaf samples were cut (4mm²) from the junction of healthy and diseased leaf tissue (Bradbury, 1975). Each sample was placed on a sterile microscopic slide, covered with a drop of sterile water and observed under a dissecting microscope (x 25). Also the sterilized a small piece of infected portion was placed in a drop of water inside a Petri-dish, teased apart and allowed to stay for 30mins to enable the multiplication of the bacterium before inoculating into the culture media.

PREPARATION OF MEDIUM AND INOCULATION OF THE PATHOGEN

The medium used in bacteria culture was nutrient agar (NA) which was prepared by weighing out 7g of readymade nutrient agar powder into a conical flask and dissolving in 250ml of sterile water (Fahy and Hayward, 1983). The mixture was thoroughly shaken and then autoclaved at 120°C for 151b for 30 minutes. This was followed by pouring 15ml of the medium into 9cm diameter Petri dishes after it has cooled down to 45°C. The nutrient agar was kept in an incubator at 28-30°C for 8 hours to enhance drying of the surface agar in the Petri dishes before use.

Prior to inoculation of bacterial suspension into the medium, the inoculation chamber was thoroughly mopped with 70% absolute alcohol using sterile cotton wool to prevent contamination. The bacterial suspension was streaked onto the nutrient agar (NA) in Petri dishes using a flamed wire loop after which the culture was placed in an incubator at 30°C for 48 hours. The culture colonies obtained after 48hrs was sub-cultured severally to obtain pure bacterial colonies. A standard bacterial inoculum concentration was obtained by serial dilution plating by addition of 5ml of sterile water into the culture colonies and then using wire loop to mix the colonies with water into suspension which was adjusted to the concentration of 10⁸ cfu/ml using haemocytometer to obtain a concentration of 10⁸ cfu/ml (colony forming unit per meal).

PATHOGENICITY TEST AND INOCULATION OF COCOYAM SEEDLINGS

The 36 pots each containing one cocoyam plant per pot were arranged in completely randomized design (CRD) with six replicates. The cocoyam seedlings were pre- inoculated using the bacterial inoculum of a concentration of 10⁸ cfu/ml before the application of plant extracts. The seedlings were inoculated by spraying the bacterial inoculum on the leaves using hand atomizer in the evening (6-6:30pm). The younger leaves and emerging shoots were also sprayed until there was a run-off. The inoculated seedlings were later covered with transparent polythene bags to create a high humid condition and allowed for 48 hours at 25-27°C for the bacterial pathogen to incubate (Jones *et al.*, 2000).

APPLICATION OF AQUEOUS PLANT EXTRACTS

The three aqueous plant extracts used were obtained from: lemon grass (*Cymbopogon citrates*) black pepper seed (*Piper guineense*), orange peel (*Citrus sinensis*). To compare their performance with standard synthetic pesticides an antibiotic (streptomycin sulphate) and a fungicide (copper oxychloride) were used as positive checks while sterile water served as untreated control. The aqueous extracts and pesticides were applied on the cocoyam leaves in the same way as the bacterial inoculation, but seven days after bacterial inoculation using an atomizer (Opara and Wokocho, 2008).

DISEASE SEVERITY INDEX AND % DISEASE INCIDENCE

Diseases severity was recorded weekly based on the scale of 0-6 as follows:

- | | | |
|---|---|---|
| 0 | = | no diseases symptom visible on the leaf surface |
| 1 | = | a few lesions but less than (<) 5% on leaf surface |
| 2 | = | 6 - 10% of the leaf surface affected by the lesions |

- 3 = 11 - 25% of the leaf surface affected
 4 = 41 - 50% or half area of the leaf surface affected
 5 = above (>) 75% of the leaf surface affected

DISEASE INCIDENCE

The disease incidence was determined as follows:

$$\text{Percentage (\%)} \text{ disease incidence} = \frac{\text{Number of leaves affected}}{\text{Total number of leaves sampled}} \times \frac{100}{1}$$

Data on growth and yield parameters were collected fortnightly, starting from one week after treatment (extract) application or four weeks after planting based on: Plant height (cm), Number of leaves, Leaf area index (cm), and the number of corms and weight of corms (g).

The scores based on first 5 leaves from the youngest open foliage were used to assess the disease severity while disease incidence was obtained by the percentage of affected number of leaves per plant and per replication.

Data Analysis

The data collected from the experiment were statistically analyzed using analysis of variance (ANOVA) and Fishers-LSD at $P < 0.05$ to separate the means according to the procedures outlined by Steel and Torrie (1980).

RESULTS

Soil Analysis and Characterization

The result of soil sample analyzed is summarized in Table1 below.

Table 1: Properties of the soil sample collected at 0–30 cm depth

Soil	Characteristics
Texture	Loamy sand
Sand (%)	61.33
Silt (%)	6.00
Clay (%)	32.66
Organic carbon (%)	1.35
organic matter(mg/kg)	15.0
Nitrogen (%)	0.11
Ca (cmo/kg ⁻¹)	2.93
Ma (cmo/kg ⁻¹)	1.26
K (cmo/kg ⁻¹)	0.16
Na (cmo/kg ⁻¹)	0.23
P (cmo/kg ⁻¹)	5.20
Ex. acidity (cmo/kg ⁻¹)	1.10
ECEC (cmo/kg ⁻¹)	5.69
pH (H2O)	4.70

Pathogenicity Test

Results of pathogenicity test conducted showed that bacterium pathogen from cocoyam induced symptoms in the leaves, which appeared within 3-4 days after inoculation and by 7-8 days, lesions increased to 1 cm in diameter, coalesced and prominent chlorotic halos developed around the lesions similar to those observed in the infected field. The control experiment however showed no such symptoms or lesions at all. Within another one week, spots and lesions on the upper surface turned gray brown with the chlorotic halos becoming enlarged, in the crops inoculated with the pathogen. These symptoms were indistinguishable from those reported for *Xanthomonas pv. disffnabachiae* in Florida (Ken *et al.*,1985).

Biochemical and Cultural Analysis of the Pathogen

The result of some biochemical and cultural tests conducted on the pathogen are shown in Tables 2 and 3 below.

Table 2: Cultural characteristics of the pathogen

Tests	Characteristics
Colour and texture of colonies	Mucoid, yellow, creamy and pale yellow
Gram staining Test	Gram negative rods
Microscopic Examination	Motile, no spores or capsules
Colony Characteristic and shape	Mucoid and convex

Table 3: Biochemical characteristics

Tests	Results
Gram reaction	-
Starch hydrolysis	+
Catalase activity	-
Gelatin hydrolysis	+
Oxidase activity	-
Acid Production from:	
Glucose	+
Mannose	+
Arabinose	+

* + = positive; - = negative

Following the results of physiological and biochemical analysis including pathogenicity tests conducted, it was concluded that the bacterium isolated from leaf spot of cocoyam in Umudike, South Eastern Nigeria for *Xanthomonas* pv. *disffenabachiae* was similar to the report made in Florida and the strain identified as *X. disffenabachiae* (Ken *et al.*, 1985).

Table 4: Effect of plant extracts on disease severity, growth parameters and yield parameters on *Colocasia* in the pot trial during the 2010 cropping season

Treatment	Disease severity	Leaf Area (cm)	No. of Leaf	Plant ht. (cm)	No. of corm	Wt. of corm /plant (kg)
Streptomycin	1.33	4.10	4.17	30.67	3.25	2.90
Copper oxychloride	1.67	3.13	3.83	30.33	3.00	2.30
<i>Cymbopogon citratus</i>	1.83	2.64	2.50	23.00	3.00	1.70
<i>Piper guineense</i>	1.33	2.95	3.50	30.00	3.60	2.60
<i>Citrus sinensis</i> peel	1.17	4.83	5.00	31.50	3.85	2.80
Sterile water control	3.00	2.14	2.00	22.17	1.00	1.30
LSD (P≤0.05)	0.57*	NS	1.81*	6.05**	1.02*	0.80*

* = significant at 5% alpha level (P≤0.05)

** = significant at 1% alpha level (P≤0.01)

NS = not significant

Effect of Plant Extracts on Disease Severity

The results obtained for the two years (2010 and 2011) on disease severity are summarized in Tables 1 and 2. Plants treated with *Citrus sinensis* peel (1.17 and 1.00) and *Piper guineense* (1.33 and 1.00) during the 2010 and 2011 trials proved more effective than the other treatments and were even as good as the standard antibiotic (streptomycin sulphate) in reducing the disease severity and their means differed significantly from that of the sterile water i.e. the control treatment (p≤0.05).

Effect of Plant Extracts on Growth Parameters

For leaf area, there was no significant differences observed in their treatments during the 2010 cropping season, however in the 2011 season plants treated with *C. sinensis* peel gave the highest leaf area (10.65) after the standard antibiotic - streptomycin (11.13cm) when compared with sterile water (7.79 cm).

Regarding the number of leaves, *C. sinensis* peel treated seedlings proved to be the highest mean number per plant (5.0) followed by streptomycin (4.17) while sterile water gave the least (2.0) which differ significantly at $P \leq 0.05$ for the two years under trials. For plant height, *C. sinensis* peel still gave the best (31.50) compare with sterile water (22.17) at $P \leq 0.05$.

Effect of Plant Extracts on Crop yield

From the results obtained during the two cropping seasons, *C. sinensis* peel produced the highest in terms of the number of corms per plant (3.85 and 8.00) followed by streptomycin (3.25 and 8.00) while the least was recorded for sterile water (1.00 and 3.00) which differ significantly from others at $P < 0.05$. However, for the weight of corm, streptomycin gave the best (2.90kg) in the first year followed by *C. sinensis* peel (2.80kg in 2010 and 2.62kg in 2011) and *P. guineense* (2.60kg for 2010 and 2.22kg in 2011), i.e. in the second year *P. guineense* (2.22kg) performed as good or even better than the standard antibiotic, streptomycin (2.08kg) in the second cropping season (i.e. 2011) next to *C. sinensis* (2.62kg) and their means differed significantly from that of control, sterile water (1.05kg) at $P \leq 0.05$.

Table 5: Effect of plant extracts on disease severity, growth parameters and yield parameters on *Colocasia* in the pot trial during the 2011 cropping season

Treatment	Disease severity	Leaf Area (cm)	No. of Leaf	Plant ht. (cm)	No. of corm	Wt. of corm /plant (kg)
Streptomycin	1.00	11.13	6.50	44.50	8.00	2.08
Copper oxychloride	1.17	7.84	6.17	45.67	7.00	1.70
<i>Cymbopogon citratus</i>	1.17	7.44	6.00	40.00	7.67	1.85
<i>Piper guineense</i>	1.00	10.26	7.83	44.00	7.00	2.22
<i>Citrus sinensis</i> peel	1.00	10.65	7.50	46.33	8.00	2.62
Sterile water control	3.49	7.79	3.00	30.00	3.00	1.05
LSD $P \leq 0.05$	0.49*	3.27*	2.12*	4.57*	2.06**	0.36**

* = significant at 5% alpha level ($P \leq 0.05$)

** = significant at 1% alpha level ($P \leq 0.01$)

Effect of Plant Extracts on Disease Incidence

Data obtained (Fig.1) during the two years trials showed that plant extracts especially *C. sinensis* (25% for both years) and *P. guineense* (25 and 28% for the 1st and 2nd year respectively) had reduced disease incidence similar to those of streptomycin which is a standard disease incidence while the untreated control recorded high disease incidence (65 and 68% for the 1st and 2nd year respectively) during the periods under investigation (Fig. 1).




Fig 1: Disease Incidence of cocoyam leaf blight during the 2010 and 2011 pot trials in Umudike.

DISCUSSION

This study shows the bactericidal potential and potency of the three plant extracts used namely *C. sinensis*, *P. guineense* and *C. citrates* in comparison with standard pesticides as far as reducing both disease severity and incidence are concerned and at the same time enhancing growth and yield of cocoyam which is an important staple food crop in south Eastern Nigeria. This work is similar to the those reported by earlier workers (Amadioha, 2003; John and James, 2004; Opara and Wokeocha, 2008) which showed that some indigenous plant species could serve as potential antimicrobial agents against bacterial pathogens. In his work Jacobson (1989) observed that some botanical pesticides were also found among the families of Asteraceae, *Lobiatae* and *Rutaceae*. From our observations, all the three plant extracts employed significantly reduced disease severity and incidence of bacterial leaf spot disease of *Colocasia* when compared with the untreated control (sterile water).

It was also noted that the plant extracts proved to be as effective as standard synthetic pesticides such as antibiotics (streptomycin sulphate) and fungicide (copper oxychloride) and were even in some cases at par with these conventional chemicals usually used in disease reduction, indicating that some natural anti-microbial active ingredients are indeed contained in these plant extracts (Stoll, 2000; John and James, 2004). Opara and Obani (2010) made similar observation in bacterial blight of egg plant. Amadioha (2004) reported that the greater effectiveness of the extracts may be due to inherent chemical constituents or bioactive ingredients of the plant extracts. These active principles in the plant extracts he concluded contain anti-bacterial polyphenolic compounds.

Accordingly the work done by Burnt (1975) showed that synthetic pesticides such as streptomycin and agromycin were found to be systemic in action similar effect could have been employed by the plant extracts on leaf blight disease of cocoyam resulting to disease reduction when compared with the untreated control. It was also noted that for the two years, foliar application of streptomycin sulphate significantly reduced disease severity three times and by 70-80% in the case disease incidence more than the control experiments. Also, other workers have observed that applying streptomycin sulphate (1000 ppm) a day prior to inoculation was more effective in the controlling bacterial diseases than after inoculation (Ken *et al.*, 1985; Whitehead, 1999). From this study it was also found that, the plant extracts significantly generally improved the amount of yield of corm and cormels further stressing the function of the bactericidal activity of the extracts used.

The specific identity of the causal agent of cocoyam leaf blight has for some time been an issue for discussion. For instance, Berniac (1974) reported that the bacterium isolated from cocoyam leaf blight was identified as *Xanthomonas campestris* pv. *dieffenbachiae* while Hayward (1983) reported that the causal agent of bacteria leaf spot of cocoyam was identified as *Xanthomonas campestris* pv. *aracearum*. These symptoms were indistinguishable from those reported by Ken *et al* (1985) for *Xanthomonas* pv. *dieffenbachiae*. On the basis of results obtained from the physiological and biochemical tests as well as the pathogenicity tests, it was concluded that the bacterium causing leaf blight of cocoyam in Umudike was a strain of *X. dieffenbachiae*, similar report was made in Florida and the strain identified as *X. dieffenbachiae* (Ken *et al.*, 1985).

Contrary to the results obtained by Berniac (1974), two of the three known *X. campestris* pv. *dieffenbachiae* isolates from other aroids grown in Florida by Ken *et al* (1985) induced typical symptoms in cocoyam but was identified as *X. campestris* pv. *dieffenbachiae* and not *X. aracearum* based on the findings that the cocoyam strains do not produce acid from glycerol. Dye and Lelliot (1974) recognized a distinct pathovar. *X. campestris* pv. *aracearum*, for the cocoyam pathogen. However, it was chosen to classify the Florida cocoyam strain as *X. campestris* pv. *dieffenbachiae* and the historical use of this pathovar designation for the xanthomonads attacking other aroids. The observations also indicate that the pathogen is systemic in cocoyam and *X. campestris* pv. *dieffenbachiae* has been considered systemic in other hosts. This is the first report of *X. campestris* pv. *dieffenbachiae* causing cocoyam leaf blight in Nigeria. The significant reduction in disease severity and incidence shows that formulations of plant extracts could have important roles in biologically based strategies for control of diseases caused by *X. campestris* pv. *dieffenbachiae*.

REFERENCES

- Agboola SD (1987). Cocoyam storage and its full potential for food sufficiency and future economic recovery of Nigeria. Proceedings of 1st National Workshop on cocoyam held at N.R.C.R.I, Umudike Nigeria, pp 42.
- Amadioha AC (2003). Evaluation of some plant leaf extracts against *Collectorichum lindemuthianum* in cowpea. Acta Phytopathol Entomol, Hungarica. 38:259-265.
- Amadioha AC (2004). Control of black rot of potato cause by *Rhizoctonia bataticola* using some plant leaf extracts. Arch. phytopathol. Plant prot. 37:111-117.
- Anikwe MAN, Mbah CN, Ezeaku PI and Onyia VN (2007). Tillage and plastic mulch effects on soil properties and growth and yield of cocoyam (*Colocasia esculenta*) on an ultisol in south-eastern Nigeria.
- Arene OB and Ene LSO (1987). Advances in cocoyam research at National Root Crops Research Institute Umudike (1972-1986). Proc. First National Workshop in Cocoyam, NRCRI, Umudike, Nigeria, pp 58-71.
- Berniac M (1974). Line Maladie bacterienne de *Xanthosoma sagittifolium* (L.) Scholt Annales de phytopathologie. 6 : 197-202.
- Bradbury JF (1970). Isolation and preliminary study of bacteria from plants. Review of Plant Pathology. 49: 213-218.
- Burnt A (1975). Effect and mode of action of some systemic nematicides. Diss. Agric. Univ. Wageningen. 3: 60-79.
- Chukwu GO and Nwosu KI (2008). Cocoyam rebirth. The renaissance of a giant crop. Paper presented at the 17th Annual conference of Nigeria Rural Sociological Association at NRCRI Umudike, 11 pp.
- Dye DW and Lelliot RA (1974). Genus *Xanthomonas* (Downson, 1939). In Bergys manual of Determinative Bacteriology, 8th edition, pp 243-249.
- Fahy PC and Hayward AC (1983). Media and culture for isolation and diagnostic tests. In: plant bacteria Diseases: a diagnostic Guide. PC Fahy and G Parsley (eds). Academic press New York, pp 15.
- Hayward AC (1983). Primary differentiation of General of plant pathogenic bacteria. In: plant bacteria Diseases: a diagnostic Guide. PC Fahy and G Parsley (eds). Academic press New York, pp 15.
- Ikwelle MG (1999). Nigeria capacities in root and tuber crops research and production. Proc. Annual review and research planting workshop NRCRI, Umudike, 23-48.
- Jacobson MO (1989). Botanical pesticides, past, present and future; insecticides of plant origin, Washington DC, USA: ACS Symposium series 387.
- John, H.B. and James C.L. (2004) Effect of formulated plant extracts and oil on population density of *Phytophthora nicotiana* in soil and control of Phytophthora blight in Greenhouse. Plant Dis. 88:11-16.
- Jones JB, Bouzar H, Stall RE, Robert PD, Bowen BW, Sunberry J, Streckler PM and Chin J (2000). Systematic analysis of (*Xanthomonas* spp.) associated with pepper and tomato lesions. International Journal of Systematic and Evolutionary Microbiology. 50:11-12.
- Ken P, Raymond BV and Wilbur D (1985). Bacteria leaf spot of cocoyam (*Xanthosoma caracu*), incited by *Xanthomonas campestris* pv. *dieffenbachiae*, in Florida.

- Opara, E.U. and Wokocha, R. C. (2008). Efficacy of some plant extract on the *in vitro* and *in vivo* control of *Xanthomonas campestris* pv. *vesicatoria*. Agriculture Journal 3(2): 163-170.
- Opara, E.U. and Obani, F.T. (2010). Performance of some plant extracts and pesticides in the control of bacterial spot diseases of *Solanum*. Agriculture Journal 5(2):45-49.
- Steel, R.G. and Torrie, J.H. (1980). Principles and Procedure of Statistics: A Biometrical Approach, 2nd ed. McGraw-Hill, New York, 633 pp.
- Stoll, G. (2000) Natural Crop Protection in the Tropics and sub tropics. Letting information come to life. 2nd edition, Margraf Veriag, pp: 102 -139.
- Whitehead, R. (1999) The Pesticides Guide British crop Protection council CABI Publishing pp. 149-159.