

[Home](#)

# *Entada africana* Guill. & Perrott.

**Leguminosae****Author: Le Houérou****Common names**

Ndende, twatsa

**Synonyms**

- *Entada sudanica* Schweinf.
- *Entada ubanguiensis* De Wild.
- *Entadopsis sudanica* (Schweinf.) Gilbert & Boutique

**Description**

Small, low branching tree, up to 7-12 m, with a narrow open crown. Bark fibrous and creviced grey-brown, transversally striped, slash red with white streaks, twigs grey-brown, pubescent. Leaves large bipinnate 15-45 cm long, with 2-9 pairs of pinnae with 8-24 pairs of linear-oblong leaflets 0.9-4.5 x 0.3-1.5 cm. Leaves glabrous. Flowers cream, in racemes 5-10 cm long. Pods, with an undulate margin, 10-30 x 2-5 cm, dark +/- pubescent, 12-15 seeds per pod, seeds two-winged, ca 4 000 per kg.

**Distribution**

South Sahel and Sudanian ecozone savannas in disturbed places, fairly common and gregarious. Throughout the Sahel to East and South Africa.

**Crop management**

Sensitive to bush fires.

**Products & uses**

Leaves are good fodder, wood soft and easy to work, Bast fibers used for bands, ropes, storage bins. Many medicinal and fetish uses : leaves; tonic tea, wound healing, contains rotenone and tannins, bark abortive, roots have antidotal effects against various toxic agents and fish poison (ichthyotoxic).

**References**

Aubréville 1950 ; Berhaut 1975 ; Hutchinson & al. 1958 ; Dalziel 1955 ; Giffard 1974a ; Kerharao & Adam 1974 ; Baumer 1975 ; Weber et al. 1977 ; Geerling 1982/88 ; Von Maydell 1983/86 ; Burkill 1995.