Descriptors for **Pistachio** (*Pistacia vera* L.)



LIST OF DESCRIPTORS

Almond (revised) * (E)	1985
Apple (E)	1982
Apricot * (E)	1984
Ávocado (E,S)	1995
Bambara groundnut (E)	1987
Banana (revised) * (E,F,S) *	1996
Barley (E)	1994
Beta (E)	1991
Black pepper (E,S)	1995
Brassica and Raphanus (E)	1990
Brassica campestris L. (E)	1987
Buckwheat (E)	1994
Capsicum (E,S)	1995
Cardamom (E)	1994
Cashew (E)	1986
Cherry * (É)	1985
Chickpea (É)	1993
Citrus (E)	1988
Coconut (E)	1992
Coffee (E.F.S) *	1996
Colocasia * (E)	1980
Cotton (Revised) (E)	1985
Cowpea (E)	1983
Cultivated potato * (E)	1977
<i>Echinochloa</i> millet * (E)	1983
Eggplant (E.F)	1990
Faba bean $*$ (E)	1985
Finger millet (E)	1985
Forage grass * (E)	1985
Forage legumes * (E)	1984
Grape * (E)	1983
Groundnut (E.F.S)	1992
Kodo millet $*$ (E)	1983
Lentil * (E)	1985
Lima bean $*$ (E)	1982
Lupin/Lupinos $*$ (E S)	1981
Maize (E, F, S)	1991
Mango (E)	1989
Medicago (Annual) * (E.F)	1991
Mung bean * (E)	1980
Oat * (E)	1985
Oca * (S)	1982
Oil palm (E)	1989
Panicum miliaceum and	
P. sumatrense (E)	1985
Papava (E)	1988
Peach * (E)	1985
Pear * (E)	1983
Pearl millet (E.F)	1993

Phaseolus acutifolius (E)	1985
Phaseolus coccineus * (E)	1983
Phaseolus vulgaris * (E)	1982
Pigeonpea (E)	1993
Pineapple (E)	1991
Pistachio (E,F) *	1997
Plum * (E)	1985
Potato variety * (E)	1985
Quinoa * (E)	1981
Rice * (E)	1980
Rye and Triticale * (E)	1985
Safflower * (E)	1983
Sesame * (E)	1981
<i>Setaria italica</i> and <i>S. pumilia</i> (E)	1985
Sorghum (E,F)	1993
Soyabean * (E, C)	1984
Strawberry (E)	1986
Sunflower * (E)	1985
Sweet potato (E,F,S)	1991
Tomato (E,F,S) *	1996
Tropical fruit * (E)	1980
Vigna aconitifolia and	
V. trilobata (E)	1985
<i>Vigna mungo</i> and	
<i>V. radiata</i> (Revised) * (E)	1985
Walnut (E)	1994
Wheat (Revised) * (E)	1985
Wheat and <i>Aegilops</i> * (E)	1978
White Clover (E)	1992
Winged bean * (E)	1979
Xanthosoma (E)	1989
Yams * (E)	1980

IPGRI publications are available free of charge to the libraries of genebanks, university departments, research institutions, etc. On request to Head, Editorial and Publications Unit, titles may also be made available to individuals who can show that they have a need for a personal copy of a publication. E, F, S and C indicate English, French, Spanish and Chinese, respectively. Titles marked * are available only as photocopies. Titles marked * are available for downloading in portable document format.

Descriptors for **Pistachio** (*Pistacia vera* L.)





The International Plant Genetic Resources Institute (IPGRI) is an autonomous international scientific organization operating under the aegis of the Consultative Group on International Agricultural Research (CGIAR). The international status of IPGRI is conferred under an Establishment Agreement which, by March 1997, had been signed by the Governments of Algeria, Australia, Belgium, Benin, Bolivia, Brazil, Burkina Faso, Cameroon, Chile, China, Congo, Costa Rica, Côte d'Ivoire, Cyprus, Czech Republic, Denmark, Ecuador, Egypt, Greece, Guinea, Hungary, India, Indonesia, Iran, Israel, Italy, Jordan, Kenya, Malaysia, Mauritania, Morocco, Pakistan, Panama, Peru, Poland, Portugal, Romania, Russia, Senegal, Slovak Republic, Sudan, Switzerland, Syria, Tunisia, Turkey, Uganda and Ukraine. IPGRI's mandate is to advance the conservation and use of plant genetic resources for the benefit of present and future generations. IPGRI works in partnership with other organizations, undertaking research, training and the provision of scientific and technical advice and information, and has a particularly strong programme link with the Food and Agriculture Organization of the United Nations. Financial support for the research agenda of IPGRI is provided by the Governments of Australia, Austria, Belgium, Canada, China, Denmark, Finland, France, Germany, India, Italy, Japan, the Republic of Korea, Luxembourg, Mexico, the Netherlands, Norway, the Philippines, Spain, Sweden, Switzerland, the UK and the USA, and by the Asian Development Bank, CTA, European Union, IDRC, IFAD, Interamerican Development Bank, UNDP and the World Bank.

Citation

IPGRI. 1997. Descriptors for Pistachio (*Pistacia vera* L.). International Plant Genetic Resources Institute, Rome, Italy.

ISBN 92-9043-332-9

This publication is available to download in portable document format from URL: http://www.cgiar.org/ipgri/

IPGRI Via delle Sette Chiese 142 00145 Rome Italy

© International Plant Genetic Resources Institute 1997

CONTENTS

PR	EFACE	iv
DE	FINITIONS AND USE OF THE DESCRIPTORS	1
PA	SSPORT	3
1.	Accession descriptors	3
2.	Collecting descriptors	5
M	ANAGEMENT	9
3.	Orchard management descriptors	9
EN	IVIRONMENT AND SITE	11
4.	Characterization and/or evaluation site descriptors	11
5.	Collecting and/or characterization/evaluation site environment descriptors	12
Cŀ	IARACTERIZATION	20
6.	Plant descriptors	20
EV	ALUATION	34
7.	Plant descriptors	34
8.	Abiotic stress susceptibility	35
9.	Biotic stress susceptibility	39
10.	Molecular markers	40
11.	Cytological characters	41
12.	Identified genes	41
RE	FERENCES	42
CC	ONTRIBUTORS	43
AC	CKNOWLEDGEMENTS	47
A٢	NEX I: List of Multi-crop Passport Descriptors	48
Ar	nex II: Collecting form for pistachio	52

PREFACE

Descriptors for Pistachio (*Pistacia vera* L.) was developed by Dr Ettore Barone, University of Palermo, Italy, Ir Paul Van Mele, University of Ghent, Belgium and Dr Stefano Padulosi, IPGRI. A draft version prepared in the internationally accepted IPGRI format for Descriptor Lists was subsequently sent to a number of international experts for their comments and amendments. A full list of the names and addresses of those involved is given in 'Contributors'.

This publication was produced within the framework of the IPGRI Project on Conservation and Use of Underutilized Mediterranean Species (UMS) (URL: http://cgiar.org/ipgri/Regional/Europe/UMS), an initiative supported by the Italian Government aiming at promoting better conservation and use of those crops indigenous to the Mediterranean region which have been neglected by science and scarcely safeguarded, in spite of their good economic potential.

IPGRI encourages the collection of data for descriptors in the first four categories of this list – *Passport, Management, Environment and Site, Characterization* – and endorses data in these categories as those that should be available for any accession. However, the number of each of the site and environment descriptor types used will depend on the crop and their importance to the crop's description. Descriptors listed under *Evaluation* allow for a more detailed description of the accession's characters, but generally require replicated site and time trials.

Although the suggested coding should not be regarded as the definitive scheme, this format represents an important tool for a standardized characterization system and it is promoted by IPGRI throughout the world.

This descriptor list is intended to be comprehensive for the descriptors that it contains. This approach assists with the standardization of descriptor definitions. IPGRI does not, however, assume that each curator will document accessions of their collection utilizing all descriptors given. Descriptors should be used when they are useful to the curator for the management and maintenance of the collection and/or to the users of the plant genetic resources. Highly discriminating descriptors are marked with stars (\star).

This descriptor list provides an international format and thereby produces a universally understood 'language' for plant genetic resources data. The adoption of this scheme for data encoding, or at least the production of a transformation method to convert other schemes into the IPGRI format, will produce a rapid, reliable and efficient means for information storage, retrieval and exchange, and will assist with the utilization of germplasm. It is recommended, therefore, that information should be produced by closely following the descriptor list with regard to: ordering and numbering descriptors, using the descriptors specified, and using the descriptor states recommended.

Annex I contains multicrop passport descriptors developed jointly by IPGRI and FAO, to provide consistent coding schemes for common passport descriptors across crops. These aim to be compatible with both future IPGRI crop descriptors lists and the FAO World Information and Early Warning System (WIEWS) on plant genetic resources.

Any suggestions for improvement on the Descriptor List for Pistachio will be highly appreciated by IPGRI.

DEFINITIONS AND USE OF THE DESCRIPTORS

IPGRI now uses the following definitions in genetic resources documentation:

Passport descriptors: These provide the basic information used for the general management of the accession (including the registration at the genebank and other identification information) and describe parameters that should be observed when the accession is originally collected.

Management descriptors: These provide the basis for the management of accessions in the genebank and assist with their multiplication and regeneration.

Environment and site descriptors: These describe the environmental and site-specific parameters that are important when characterization and evaluation trials are held. They can be important for the interpretation of the results of those trials. Site descriptors for germplasm collecting are also included here.

Characterization descriptors: These enable an easy and quick discrimination between phenotypes. They are generally highly heritable, can be easily seen by the eye and are equally expressed in all environments. In addition, these may include a limited number of additional traits thought desirable by a consensus of users of the particular crop.

Evaluation descriptors: Many of the descriptors in this category are susceptible to environmental differences but are generally useful in crop improvement and others may involve complex biochemical or molecular characterization. They include yield, agronomic performance, stress susceptibilities and biochemical and cytological traits.

Characterization will normally be the responsibility of genebank curators, while evaluation will typically be carried out elsewhere (possibly by a multidisciplinary team of scientists). The evaluation data should be fed back to the genebank which will maintain a data file.

Highly discriminating descriptors in this descriptor list are marked with stars (\bigstar).

The following internationally accepted norms for the scoring, coding and recording of descriptor states should be followed:

- (a) the Système International d'Unités (SI units) is used;
- (b) the units to be applied are given in square brackets following the descriptor name;
- (c) standard colour charts, e.g. Royal Horticultural Society Colour Chart, Methuen Handbook of Colour, or Munsell Color Chart for Plant Tissues, are strongly recommended for all ungraded colour characters (the precise chart used should be specified in the section where it is used);

3

- (d) many quantitative characters which are continuously variable are recorded on a 1-9 scale, where:
 - 1Very low6Intermediate to high2Very low to low7High
 - Low 8 High to very high
 - 4 Low to intermediate 9
 - 5 Intermediate

is the expression of a character. The authors of this list have sometimes described only a selection of the states, e.g. 3, 5 and 7 for such descriptors. Where this has occurred, the full range of codes is available for use by extension of the codes given or by interpolation between them, e.g. in Section 9 (Biotic stress susceptibility) 1 = very low susceptibility and 9 = very high susceptibility;

Very high

(e) when a descriptor is scored using a 1-9 scale, such as in (c), '0' would be scored when (i) the character is not expressed; (ii) when a descriptor is not applicable. In the following example, '0' will be recorded if an accession does not have a central leaf lobe:

Shape of central leaf lobe

- 3 Toothed
- 5 Elliptic
- 7 Linear
- (f) absence/presence of characters is scored as in the following example:

Absence/presence of terminal leaflet

0	Absent	
1 (or +)	Present	

- (g) blanks are used for information not yet available;
- (h) for accessions which are not generally uniform for a descriptor (e.g. mixed collection, genetic segregation), the mean and standard deviation could be reported where the descriptor is continuous. Where the descriptor is discontinuous, several codes in the order of frequency could be recorded; or other publicized methods can be utilized, such as R.S. Rana *et al.* (1991) or van Hintum (1993), that clearly state a method for scoring heterogeneous accessions;
- (i) dates should be expressed numerically in the format YYYYMMDD, where

YYYY	-	4 digits to represent the year
MM	-	2 digits to represent the month
DD	-	2 digits to represent the day.

PASSPORT

1. Accession descriptors

★ 1.1 Accession number

This number serves as a unique identifier for accessions and is assigned when an accession is entered into the collection. Once assigned this number should never be reassigned to another accession in the collection. Even if an accession is lost, its assigned number should never be re-used. Letters should be used before the number to identify the genebank or national system (e.g. IDG indicates an accession that comes from the genebank at Bari, Italy; CGN indicates an accession from the genebank at Wageningen, The Netherlands; PI indicates an accession within the USA system).

1.1.1 Local plant number

This identifies a single plant within a population of plants having the same accession number. It may be any combination of plot identity, row number, or tree position within the row

1.2 Donor name

Name of institution or individual responsible for donating the germplasm

1.3 Donor number

Number assigned to an accession by the donor

1.4 Country where maintained

Name of the country in which the sample is maintained. Use the three-letter abbreviations from the *International Standard (ISO) Codes for the representation of names of countries*, No. 3166, 4th Edition. Copies of these are available from DIN: Deutsche Institut für Normung e.V., D-10772 Berlin, Germany; Tel. 30-2601-2860; Fax 30-2601-1231, Tlx. 184 273-din-d.

1.5 Site where maintained

Name of institution in which collection is maintained

1.6 Curator's name

Name of officer responsible for maintaining the genetic resources material held at the site specified in descriptor **1.5 Site where maintained**

1.7 Other number(s) associated with the accession

Any other identification number known to exist in other collections for this accession, e.g. USDA Plant Inventory number (not Collecting number, see descriptor **2.3**). Other numbers can be added as 1.4.3, etc.

- 1.7.1 Other number 1
- 1.7.2 Other number 2

1.8 Scientific name

- ★ 1.8.1 Genus
- ★ 1.8.2 Species
- ★ 1.8.3 Subspecies
 - 1.8.4 Botanical variety

1.9 Genetic origin

- 1 Open pollination
- 2 Artificial pollination
- 3 Clonal selection

1.10 Pedigree

Parentage or nomenclature, and designations assigned to breeders' material. For interspecific hybrids, the species should be designated as 'hybrid' and the parentage indicated here

★ 1.11 Sex

Male 2 Female

1.12 Accession

1

1.12.1 Accession name

Either a registered or other formal designation given to the accession

1.12.2 Local language

Language in which the accession name is given

1.12.3 Translation/Transliteration

Provide translation of the local cultivar name into English

1.12.4 Year of release of the accession/year of registration

1.12.5 Synonyms

Include here any previous identification other than the current name. Collecting number or newly assigned station name are frequently used as identifiers.

1.13 Acquisition date [YYYYMMDD]

Date on which the accession entered the collection

★ 1.14 Type of material received

Seed

- *In vitro* plant 4 Bud
- Cutting 99 Other (e.g. more than one type,
 - specify in descriptor **1.16** Notes)

1.15 Accession size

Number of trees/shrubs of an accession or approximate number of seeds (if artificially pollinated) of an accession in the genebank

1.16 Notes

1

2

3

Any additional information may be specified here

2. Collecting descriptors

★ 2.1 Collecting institute(s)

Institute(s) and people collecting/sponsoring the sample collection

2.2 Site number

Number assigned to the physical site by the collector

★ 2.3 Collecting number

Original number assigned by the collector(s) of the sample, normally composed of the name or initials of the collector(s) followed by a number. This item is essential for identifying duplicates held in different collections. It should be unique and always accompany subsamples wherever they are sent.

2.4 Collecting date of original sample [YYYYMMDD]

2.5 Country of collecting

Name of the country in which the sample was collected. Use the three-letter abbreviations from the *International Standard (ISO) Codes for the representation of names of countries*, No. 3166, 4th Edition. Copies of these are available from DIN: Deutsche Institut für Normung e.V., D-10772 Berlin, Germany; Tel. 30-2601-2860; Fax 30-2601-1231, Tlx. 184 273-din-d.

2.6 Province/State

Name of the primary administrative subdivision of the country in which the sample was collected

2.7 Department/County

Name of the secondary administrative subdivision (within a Province/State) of the country in which the sample was collected

2.8 Location of collecting site

Distance in kilometers and direction from the nearest town, village or map grid reference point (e.g. CURITIBA 7S means 7 km south of Curitiba)

2.9 Latitude of collecting site

Degrees and minutes followed by N (North) or S (South) (e.g. 1030S). Missing data (minutes) should be indicated with hyphen (e.g. 10—S).

2.10 Longitude of collecting site

Degrees and minutes followed by E (East) or W (West) (e.g. 07625W). Missing data (minutes) should be indicated with hyphen (e.g. 076—W).

2.11 Elevation of collecting site [m asl]

2.12 Collecting source

- 0 Unknown
- 1 Wild habitat
 - 1.1 Forest/woodland
 - 1.2 Shrubland
 - 1.3 Grasslands
 - 1.4 Desert/tundra
- 2 Farm
 - 2.1 Field
 - 2.2 Orchard
 - 2.3 Garden
 - 2.4 Fallow
 - 2.5 Pasture
 - 2.6 Store
- 3 Market
 - 3.1 Town
 - 3.2 Village
 - 3.3 Urban area (around city)
 - 3.4 Other exchange system
- 4 Institute/Research organization
- 99 Other (specify in descriptor 2.26 Collector's notes)

★ 2.13 Number of samples collected

★ 2.14 Type of sample

Form of sample collected. If different types of material were collected from the same source, each sample type should be designated with a unique collecting number and a corresponding unique accession number

- 1 Vegetative
- 2 Seed
- 3 Pollen
- 4 Tissue culture

2.15 Status of sample

- 0 Unknown
- 1 Wild
- 2 Weedy
- 3 Traditional cultivar/Landrace
- 4 Breeders line
- 5 Advanced cultivar
- 99 Other (specify in descriptor 2.26 Collector's notes)

2.16 Uses of the accession

- 1 Nut production
- 2 Clonal rootstock
- 3 Seedling rootstock
- 4 Pollinator
- 5 Medicinal
- 6 Forage
- 7 Wood/timber
- 99 Other (specify in descriptor 2.26 Collector's notes)

2.17 Ethnic group

Name of the ethnic group of the farmer donating the sample or of the people living in the area of collecting

2.18 Local/vernacular name

Name given by farmer to crop and cultivar/landrace/weed. State language and dialect if the ethnic group is not provided

2.19 Collecting site population structure

2.19.1 Number of trees sampled

2.19.2 Frequency of accession at collecting site

- 1 Rare
- 3 Occasional
- 5 Frequent
- 7 Abundant
- 9 Very abundant

2.19.3 Associated flora

Other dominant crop/plant species, found in and around the collecting site

2.19.4 Associated mycorrhizal fungi

Were root samples collected? If so, specify which fungi were identified in the laboratory in descriptor **2.26 Collector's notes.**

- 0 No
- 1 Yes

2.20 Herbarium specimen

Was a herbarium specimen collected? If so, provide an identification number and indicate in which place (herbarium) the pistachio specimen was deposited, in descriptor **2.26 Collector's notes.**

- 0 No
- 1 Yes

2.21 Photograph

Was a photograph(s) taken of the accession or habitat at the time of collecting? If so, provide an identification number(s) in descriptor **2.26 Collector's notes**.

- 0 No
- 1 Yes

2.22 Collecting source environment

Use descriptors **5.1.1** to **5.1.22** in section 5

2.23 Cultural methods

2.23.1 Cropping system

- 1 Monoculture (specify spacing)
- 2 Intercropping (specify spacings and type of intercrop)
- 3 Agropastoralism (specify type of animals)
- 4 Natural cropping (i.e. wild *Pistacia* species topworked with cultivar)

2.23.2 Propagation method

Method used to produce trees

- 1 Seed
- 2 Grafted (specify species, hybrid and/or clone used as rootstock)
- 3 Tissue culture

2.23.3 Irrigation

- 1 Rainfed
- 2 Irrigated (specify average annual amount of water supplied per hectare)
- 3 Run off
- 4 River banks
- 99 Other (specify in descriptor **2.26 Collector's notes**)

2.23.4 Pruning

- 1 Light (<20% of the whole plant)
- 2 Medium (20 to 40%)
- 3 Severe (>40%)

2.24 Plant population density

Quantify plants by hectare

2.25 Prevailing stresses

Information on associated biotic and abiotic stresses and the accession's reaction. Indicate stresses in descriptor **2.26 Collector's notes**.

2.26 Collector's notes

Additional information recorded by the collector (e.g. assessment of genetic erosion) or any specific information on any state in any of the above descriptors

MANAGEMENT

3 Orchard management descriptors

3.1 Accession number

(Passport 1.1)

3.1.1 Local plant number (Passport 1.1.1) This identifies a single plant within a population of plants having the same accession number. It may be any combination of plot identity, row number, or tree position within the row

3.2 Accession orchard location

Enter separate block designations, row numbers and tree numbers within the row for each duplicate tree of each accession if each tree is not identified with a unique local plant number (see descriptor **3.1.1**)

- 3.2.1 Block designation
- 3.2.2 Row number
- 3.2.3 Tree number within the row

3.3 Propagation method

Method used to produce trees

- 1 Seed
- 2 Grafted (specify method used in descriptor **3.11 Notes**)
- 3 Tissue culture

3.4 Rootstock

Indicate the name of the rootstock used in descriptor 3.11 Notes

3.5 Grafting establishment [%]

Percentage of grafts that were successful

3.6 Planting year [YYYY]

Specify year tree was planted in the orchard

3.7 Regeneration year [YYYY]

Year (estimate) tree should be propagated for regeneration

3.8 Date of last regeneration or multiplication [YYYYMMDD]

Primary method of regeneration is propagation of clonal material

3.9 Number of times accession regenerated

Since the date of acquisition

★ 3.10 Type of maintenance

- 1 Vegetative in the field
- 2 Vegetative in tissue culture
- 3 Pollen
- 4 Seed
- 99 Other (e.g. more than one type, specify in descriptor **3.11 Notes**)

3.11 Notes

Any additional information may be specified here

ENVIRONMENT AND SITE

4. Characterization and/or evaluation site descriptors

4.1 Country of characterization and/or evaluation

(See instructions in descriptor 2.5 Country of collecting)

4.2 Site (research institute)

4.2.1 Latitude

Degrees and minutes followed by N (North) or S (South) (e.g. 1030S). Missing data (minutes) should be indicated with hyphen (e.g. 10—S).

4.2.2 Longitude

Degrees and minutes followed by E (East) or W (West) (e.g. 07625 W). Missing data (minutes) should be indicated with hyphen (e.g. 076—W).

- 4.2.3 Elevation [m asl]
- 4.2.4 Name of farm or institute
- 4.3 Evaluator's name and address
- 4.4 Sowing or grafting date [YYYYMMDD]

4.5 Evaluation environment

Environment in which characterization/evaluation was carried out

- 1 Field
- 2 Screenhouse
- 3 Glasshouse
- 4 Laboratory
- 99 Other (specify in descriptor 4.14 Notes)

4.6 Condition of tree

4

Choose the one condition that best fits the accession at the time of characterization/evaluation

- 1 Dying 5 Mature vigorous
 - 6 Young (not yet bearing)
- 2 Old declining
 3 Mature diseased
- 7 Healthy cropping poorly
- Mature non-vigorous 8 Healthy cropping well

4.7 Seed germination [%]

Specify number of days over which germination is measured

4.8 Field establishment [%]

Specify number of days over which establishment is measured

4.9 Sowing site in the field

Give block, strip and/or row/plot numbers as applicable, plants/plot, replication

4.10 **Field spacing**

4.10.1 Distance between trees in a row [m]

4.10.2 Distance between rows [m]

4.11 Fertilizer

Specify types, doses, frequency of each and method of application

Plant protection 4.12

Specify pesticides used, doses, frequency of each and method of application

Environmental characteristics of site 4.13

Use descriptors 5.1.1 to 5.1.22 in section 5

4.14 Notes

Any other site-specific information

5. Collecting and/or characterization/evaluation site environment descriptors

5.1 Site environment

\star 5.1.1 Topography

This refers to the profile in elevation of the land surface on a broad scale. The reference is FAO (1990)

1	Flat	0 - 0.5%
2	Almost flat	0.6 - 2.9%
3	Gently undulating	3 - 5.9%
4	Undulating	6 - 10.9%
5	Rolling	11 - 15.9%
6	Hilly	16 - 30%
7	Steeply dissected	>30%, moderate elevation range
8	Mountainous	>30%, great elevation range (>300 m)
99	Other	(specify in appropriate section's Notes)

\star 5.1.2 Higher level landform (general physiographic features)

The landform refers to the shape of the land surface in the area in which the site is located (adapted from FAO 1990)

- 1 Plain
- 5 Upland
- 2 Basin 3
- Hill
- Valley
- 6 7 Mountain
- 4 Plateau

5.1.3 Land element and position

Description of the geomorphology of the immediate surroundings of the site (adapted from FAO 1990). (See Fig. 1)

- 1 Plain level
- 2 Escarpment
- 3 Interfluve
- 4 Valley
- 5 Valley floor
- 6 Channel
- 7 Levee
- 8 Terrace
- 9 Floodplain
- 10 Lagoon
- 11 Pan

25

19

26

2

- 12 Caldera
- 13 Open depression

14

14 Closed depression

13

- 15 Dune
- 16 Longitudinal dune
- 17 Interdunal depression
- 18 Mangrove
- 19 Upper slope
- 20 Midslope
- 21 Lower slope
- 22 Ridge
- 23 Beach
- 24 Beachridge
- 25 Rounded summit
- 26 Summit
- 27 Coral atoll
- 28 Drainage line (bottom position in flat or almost-flat terrain)
- 29 Coral reef
- 99 Other (specify in appropriate section's **Notes**)



★ 5.1.4 Slope [°]

Estimated slope of the site

5.1.5 Slope aspect

The direction that the slope on which the accession was collected faces. Describe the direction with symbols N, S, E, W (e.g. a slope that faces a southwestern direction has an aspect of SW)

5.1.6 **Crop agriculture**

(Adapted from FAO 1990)

5.1.6.1 Tree and shrub cropping

- Non-irrigated tree crop cultivation 1
- 2 Irrigated tree crop cultivation
- Non-irrigated shrub crop cultivation 3
- 4 Irrigated shrub crop cultivation

5.1.7 Overall vegetation surrounding and at the site

(Adapted from FAO 1990)

- 1 Grassland (Grasses, subordinate forbs, no woody species)
- 2 Forbland (Herbaceous plants predominant)
- 3 Forest (Continuous tree layer, crowns overlapping, large number of tree and shrub species in distinct layers) 4 Woodland (Continuous tree layer, crowns usually not touching,
- understorey may be present)
- 5 Shrubland (Continuous layer of shrubs, crowns touching)
- 6 Savanna (Grasses with a discontinuous layer of trees or shrubs)
- 99 Other (specify in appropriate section's Notes)

5.1.8 Soil parent material

(Adapted from FAO 1990)

Two lists of examples of parent material and rock are given below. The reliability of the geological information and the knowledge of the local lithology will determine whether a general or a specific definition of the parent material can be given. Saprolite is used if the in situ weathered material is thoroughly decomposed, clayrich but still showing rock structure. Alluvial deposits and colluvium derived from a single rock type may be further specified by that rock type.

5.1.8.1 **Unconsolidated material**

2

4

- 1 Aeolian deposits
- (unspecified)
- 6 Lacustrine deposits
- 7 Fluvial deposits
- 8 Alluvial deposits
- Aeolian sand 3 Littoral deposits Lagoonal deposits
- 9 Unconsolidated (unspecified)
- 5 Marine deposits
- 10 Volcanic ash

- 11 Loess
- 12 Pyroclastic deposits
- 13 Glacial deposits
- 14 Organic deposits
- 15 Colluvial deposits

5.1.8.2 Rock type

(Adapted from FAO 1990)

- 1 Acid igneous/
- metamorphic rock
- 2 Granite
- 3 Gneiss
- 4 Granite/gneiss
- 5 Quartzite
- 6 Schist
- 7 Andesite
- 8 Diorite
- 9 Basic igneous/ metamorphic rock
- 10 Ultra basic rock
- 11 Gabbro
- 12 Basalt
- 13 Dolerite
- 14 Volcanic rock
- 15 Sedimentary rock

- 16 In situ weathered
- 17 Saprolite
- 99 Other (specify in appropriate section's **Notes**)
- 16 Limestone
- 17 Dolomite
- 18 Sandstone
- 19 Quartzitic sandstone
- 20 Shale
- 21 Marl
- 22 Travertine
- 23 Conglomerate
- 24 Siltstone
- 25 Tuff
- 26 Pyroclastic rock
- 27 Evaporite
- 28 Gypsum rock
- 99 Other (specify in appropriate
 - section's Notes)
- 0 Not known

5.1.9 Stoniness/rockiness/hardpan/cementation

- 1 Tillage unaffected
- 2 Tillage affected
- 3 Tillage difficult
- 4 Tillage impossible
- 5 Essentially paved

5.1.10 Soil drainage

(Adapted from FAO 1990)

- 3 Poorly drained
- 5 Moderately drained
- 7 Well drained

5.1.11 Soil salinity

- 1 <160 ppm dissolved salts
- 2 160 240 ppm
- 3 241 480 ppm
- 4 >480 ppm

5.1.12 Soil depth to groundwater table

(Adapted from FAO 1990)

The depth to the groundwater table, if present, as well as an estimate of the approximate annual fluctuation, should be given. The maximum rise of the groundwater table can be inferred approximately from changes in profile colour in many, but not all, soils.

- 0 25 cm 1
- 2 25.1 - 50 cm
- 3 50.1 - 100 cm
- 100.1 150 cm 4
- 5 >150 cm

5.1.13 Soil matrix colour

(Adapted from FAO 1990)

The colour of the soil matrix material in the root zone around the accession is recorded in the moist condition (or both dry and moist condition, if possible) using the notation for hue, value and chroma as given in the Munsell Soil Color Charts (Munsell Color 1977). If there is no dominant soil matrix colour, the horizon is described as mottled and two or more colours are given and should be registered under uniform conditions. Early morning and late evening readings are not accurate. Provide depth of measurement (cm). If colour chart is not available, the following states may be used:

- White 1
- 8 Yellowish brown
- 15 Bluish-black 16 Black

- 2 Red 3 Reddish
 - 10 Reddish yellow
 - 11 Greenish, green

Yellow

- Yellowish red Brown
- 5 6 Brownish
 - 13 Greyish **Reddish brown**
 - 14 Blue

9

5.1.14 Soil pH

4

7

Actual value of the soil within the following root depths around the accession

12 Grey

- 5.1.14.1 pH at 10-15 cm
- 5.1.14.2 pH at 16-30 cm
- 5.1.14.3 pH at 31-60 cm
- 5.1.14.4 pH at 61-90 cm
- \star 5.1.15 Soil erosion
 - 3 Low
 - Intermediate 5
 - 7 High

5.1.16 Rock fragments

(Adapted from FAO 1990)

1

Large rock and mineral fragments (>2 mm) are described according to abundance

- 0 2% 4
- 2.1 5%
 3 5.1 15%
- 5 40.1 80%

15.1 - 40%

6 >80%

★ 5.1.17 Soil texture classes

(Adapted from FAO 1990)

For convenience in determining the texture classes of the following list, particle size classes are given for each of the fine earth fractions below. (See Fig. 2)

- 1 Clay
- 2 Loam
- 3 Clay loam
- 4 Silt
- 5 Silty clay
- 6 Silty clay loam
- 7 Silt loam
- 8 Sandy clay
- 9 Sandy clay loam
- 10 Sandy loam
- 11 Fine sandy loam

- 12 Coarse sandy loam
- 13 Loamy sand
- 14 Loamy very fine sand
- 15 Loamy fine sand
- 16 Loamy coarse sand
- 17 Very fine sand
- 18 Fine sand
- 19 Medium sand
- 20 Coarse sand
- 21 Sand, unsorted
- 22 Sand, unspecified



Fig. 2 Soil texture classes

5.1.17.1 Soil particle size classes

(Adapted from FAO 1990)

1	Clay			< 2 µm
2	Fine silt	2	-	20 µm
3	Coarse silt	21	-	63 µmm
4	Very fine sand	64	-	125 µm
5	Fine sand	126	-	200 µm
6	Medium sand	201	-	630 µm
7	Coarse sand	631	-	1250 µm
8	Very coarse sand	1251	-	2000 µm

★ 5.1.18 Soil taxonomic classification

As detailed a classification as possible should be given. This may be taken from a soil survey map. State class (e.g. Alfisols, Spodosols, Vertisols, etc.).

★ 5.1.19 Water availability

- 1 Rainfed
- 2 Irrigated
- 3 Flooded
- 4 River banks
- 5 Sea coast
- 99 Other (specify in appropriate section's **Notes**)

★ 5.1.20 Soil fertility

General assessment of the soil fertility based on existing vegetation

- 3 Low
- 5 Moderate
- 7 High

★ 5.1.21 Climate of the site

Should be assessed as close to the site as possible

★ 5.1.21.1 Temperature [°C]

Provide either the diurnal (mean, maximum, minimum) or the seasonal (mean, maximum, minimum)

 ★ 5.1.21.2 Rainfall [mm] Annual average (state number of recorded years)

5.1.21.3	Wind [km/s]]		
Annual	Annual average (state number of years recorded)			
	5.1.21.3.1	Frequency of typhoons or hurricane force windsLowIntermediateHigh		
	5.1.21.3.2	Date of most recent typhoons or hurricane force winds [YYYYMMDD]		
	5.1.21.3.3	Annual maximum wind velocity [km/s]		
5.1.21.4	Frost			
	5.1.21.4.1	Date of most recent frost [YYYYMMDD]		
	5.1.21.4.2	Minimum temperature [°C]		
	Specify sea	sonal average and minimum survived		
	5.1.21.4.3	Duration of temperature below 0°C [d]		
5.1.21.5	Relative hu	midity		
	5.1.21.5.1	Relative humidity diurnal range [%]		
	5.1.21.5.2	Relative humidity seasonal range [%]		
5.1.21.6	Light 3 Shady 7 Sunny			

5.1.22 Other

(Specify in appropriate section's Notes)

CHARACTERIZATION

6. Plant descriptors

6.1.1

Average of at least two 'on-years' (production years) data, unless otherwise stated

6.1 **Growth descriptors** Reference variety

Tree vigour Bianca, Kirmizi, M-57, Sfax, 02-18 5 Intermediate Aegina, Alpha, Kerman, M-502 Ajami, Beta, Boundoky, Marawhy, Mateur, M-37, Ouleimy, Red Aleppo, Siirt

6.1.2 **Growth habit**

- (See Fig. 3)
 - 1 Erect

3 Low

7 High

- 2 Semi-erect
- 3 Spreading
- Drooping 4

Ashoury, Larnaka, Maknassy Alpha, Kerman, Sfax, Uzun Aegina, Mateur, Cerasola, Djalab, Ahmar, Gamma, Oady Batoury, Eirnora, Joley





Fig. 3 Growth habit

6.1.3 **Branching habit**

7

5

7

3	Sparse	Ashoury, Gamma, Larnaka
5	Intermediate	Kerman, Ouleimy

Alpha, Beta, Marawhy Dense

6.1.4 **Apical dominance**

To be estimated as number of lateral branches on one- and two-year old wood

- Weak 3
 - Marawhy Intermediate Batoury, Kerman
 - Cerasola, Gamma, Larnaka Strong

6.2 Leaf descriptors

For the following descriptors, average of 20 fully expanded representative leaves, collected from different trees when shoots are lignified. Do not select leaves that are out of the ordinary due to disease, nutritional imbalances and excessive vigour. For qualitative characteristics, indicate the predominant one

6.2.1 Leaf length [cm]

Measured from the base of petiole to the tip of terminal leaflet. (See Fig. 4)

6.2.2 Leaf width [cm]

Measured at the widest part. (See Fig. 4)



Fig. 4 Leaf length and width

6.2.3 Number of leaflets

- 6.2.4 Terminal leaflet length [cm] ★ (See Fig. 5)
- 6.2.5 Terminal leaflet width [cm] ★ Measured at the widest part. (See Fig. 5)



Fig. 5 Terminal leaflet length and width

★ 6.2.6 Terminal leaflet/width ratio

6.2.7 Terminal leaflet size

- 1 Similar to basal leaflets
- 2 Larger than basal leaflets

6.2.8 Terminal leaflet shape

(See Fig. 6)

- 1 Broad lanceolate
- 2 Elliptic
- 3 Ovate
- 4 Round ovate
- 5 Roundish
- 99 Other (specify in descriptor 6.6 Notes)



Fig. 6 Terminal leaflet shape

6.2.9 Terminal leaflet apex

(See Fig. 7)

- 1 Acuminate
- 2 Mucronate
- 3 Mucronulate
- 4 Obtuse
- 5 Retuse
- 99 Other (specify in descriptor 6.6 Notes)



Fig. 7 Terminal leaflet apex

6.2.10 Terminal leaflet base

(See Fig. 8)

- 1 Attenuate
- 2 Obtuse
- 3 Truncate
- 4 Oblique
- 99 Other (specify in descriptor 6.6 Notes)



Fig. 8 Terminal leaflet base

- 6.2.11 Terminal leaflet margin
 - 1 Flat
 - 2 Wavy

6.2.12 Petiole shape

- 1 Flattened
- 2 Rounded
- 3 Rounded straight adaxially

6.2.13 Leaf colour

Evaluated at adaxial side, at time when shoot is woody, before harvest time

- Light green 1
- 2 Green
- 3 Dark green

6.2.14 One-year-old shoot colour

To be evaluated at winter time

- 1 Very light brown
- 2 Light brown
- 3 Brown

6.2.15 Leaf midrib indumentum

- Glabrous 1
- 2 Sparsely puberulent

6.3 Inflorescence and fruiting habit

Average over at least two 'on-years' (except for descriptors 6.3.4 and 6.3.5 for females). Bud descriptors are evaluated at harvest time, inflorescence descriptors at peak bloom period

\star 6.3.1 Female reference standard

Indicate which cultivar has been used for the following descriptors where applicable. If possible use one of the listed cultivars. If not available, use main locally used cultivar

1	Aegina	7	Mateur
2	Ashoury	8	Ohadi
3	Batoury	9	Sfax
4	Bianca	10	Siirt
5	Kerman	99	Other (specify in
6	Larnaka		the descriptor 6.6 Notes)

6.3.2 Male reference standard \star

Indicate which cultivar has been used for the following descriptors where applicable. If possible use one of the listed cultivars. If not available, use main locally used cultivar 7

- 1 Alpha (syn. A) 2 Ask
- 8 Nazareth (syn. Naz.)

M-57

M-11

10 Peters

- Beta (syn B)
- 4 Chico
- 5 Enk

9

- Gamma
- 99 Other (specify in the descriptor 6.6 Notes)

\star 6.3.3 **Flowering precocity**

6

3

Specify number of years from Graft or Seed to first flower (i.e. 4 G indicates first flower produced 4 years from graft establishment)

6.3.3.1 Years before (-) or after (+) reference standard6.3.3.2 Years from graft or seed to first yield

Of at least 300 nuts/tree. Specify number of years as above

6.3.4 Inflorescence bud dry weight [DW g]

Average of 20 buds during off-years for female¹

6.3.5 Inflorescence bud shape

 \star

Evaluated during off-years for female. (See Fig. 9)

- 1 Broadly ovate
- 2 Narrowly ovate
- 3 Conical



Fig. 9 Inflorescence bud shape

6.3.6 Inflorescence bud colour

- 1 Reddish brown
- 2 Light brown
- 3 Brown
- 4 Dark brown

6.3.7 Inflorescence abundance

Rate in relation to reference standard of same age

- 3 Sparse
- 5 Intermediate
- 7 Dense

6.3.8 Inflorescence rachis length [cm]

Average of 20 inflorescences at peak bloom period

Weight of buds here as well as the weight of kernels and nuts in following descriptors should always be calculated using material dried in a ventilated oven at 60°C for 24 h

6.3.9 Number of primary lateral inflorescence branches

Average of 20 inflorescences at peak bloom period

6.3.10 Alternate bearing

2

Estimated as percentage of inflorescence bud drop in on-years

- 1 Slight < 35%
 - Moderate 35% 65%
 - 3 Significant > 65%

6.4 Nut and kernel

Descriptors within this section should be used on healthy nuts at harvest time, unless otherwise specified. (See Fig. 10)



Fig. 10 Lateral section of a Pistacia vera fruit

6.4.2 Hull consistency

- 1 Juicy
- 2 Dry

6.4.3 Hull tip

Evaluated at maturity

- 3 Little pronounced
- 5 Pronounced
- 7 Strongly pronounced

6.4.4 Hull colour

- 1 Light cream
- 1 Yellow-white group
- 2 Orange-white group
- 3 Yellow-orange group
- 4 Orange-red group
- 5 Red group
- 6 Red-purple group
- 99 Other (specify in descriptor **6.6 Notes**)

6.4.5 Hull colour homogenity

(See Fig. 11)

- 0 No (Hull tip colour clearly different from rest of hull) Batoury
- 1 Yes (Colour equally distributed)



Ashoury

Fig. 11 Hull colour homogeneity

★ 6.4.6 Nut length [mm]

Average of 20 nuts, measured from most distant points along main seed axis. (See Fig. 12)

★ 6.4.7 Nut width [mm]

Average of 20 nuts, measured from the widest points perpendicular to main seed axis (See Fig. 12)

★ 6.4.8 Nut thickness [mm]

Average of 20 nuts, measured at widest part perpendicular to suture. (See Fig. 12)



Fig. 12 Nut length, width and thickness

Ghochi, Kaleh Kerman, Ohadi

Nab-al-Djamal, Joley, Uzun

★ 6.4.9 Nut shape

(See Fig. 13)

- 1 Roundish (l/w < 1.5)
- 2 Ovoid (1.5 < l/w < 1.8) Batoury, Bianca, Mateur, Red Jalap
- 3 Elongated (l/w > 1.8)
- 4 Narrowly cordate
- 5 Cordate





Fig. 13 Nut shape

★ 6.4.10 Shell apex

(See Fig. 14)

- /		
1	Flattened	Batoury
2	Rounded	Uzun
3	Symmetrically pointed	Marawhy
4	Asymmetrically pointed	Mateur



6.4.12 Nut abscission

- 3 Easy
- 5 Medium
- 7 Difficult

6.4.13 Pedicel scar colour

- 1 Lighter than shell colour
- 2 Similar to shell colour
- 3 Darker than shell colour

Pedicel scar shape 1 Orbicular

- 2 Ovate
- 3 Elliptic
- 4 Elongated
- 99 Other (specify in descriptor 6.6 Notes)

Siirt, Nab-al-Djamal Uzun





3 Low 7 High Marawhy, Mateur Batoury





★ 6.4.17 Split nuts [%]

★ 6.4.18 Tendency to early nut splitting

Measured four weeks prior to harvest

- Low Red Aleppo
- Moderate Kerman
- 7 High

3

5

 \star

6.4.19

Position of suture opening

- 1 Dorsal side only
- 2 Mainly dorsal side
- 3 Ventral side only
- 4 Mainly ventral side
- 5 Dorsal and ventral side completely

★ 6.4.20 Suture opening

Evaluated at harvest time. (See Fig. 18)

- 3 Narrow
- 5 Moderate
- 7 Wide

Uzun Ashoury Ohari, Kaleh Ghochy



Fig. 18 Suture opening

6.4.21 Tendency to shell staining

Record at har	vest time	
3	Low	Uzun
5	Intermediate	Ashoury
7	High	Mateur, Nab-al-Djamal

- ★ 6.4.22 Blank production [%] (Approximate percentage)
- ★ 6.4.23 100-Nut weight [DW g] Record the mean of healthy dry nuts
- ★ 6.4.24 Number of nuts in 100 g Evaluated using healthy ready-for storage nuts
- ★ 6.4.25 100-Kernel weight [g] Average of healthy dry kernels

★ 6.4.26 Kernel dry weight/nut dry weight X 100

6.4.27 Kernel length [mm]

Average of 20 kernels, measured from most distant points along main seed axis. (See Fig. 19)

6.4.28 Kernel width [mm]

Average of 20 kernels, measured on the widest points perpendicular to main seed axis (See Fig. 19)

6.4.29 Kernel thickness [mm]

Average of 20 nuts, measured at the widest part perpendicular to cotyledon suture. (See Fig. 19)



Fig. 19 Length, width and thickness of kernel

6.4.30 Kernel flavour

- 1 Satisfactory
- 2 Unsatisfactory

★ 6.4.31 Kernel colour

Based on 20 randomly selected kernels.

- 1 Yellowish Kerman
- 2 Yellowish green Ashoury, Mateur

Bianca

- 3 Green
- 99 Other (specify in descriptor 6.6 Notes)

★ 6.4.32 Testa colour

Based on 20 randomly selected kernels.

1	Greyish	Boundouky
2	Reddish	Oleimy
3	Deep red	Bayadi

99 Other (specify in descriptor 6.6 Notes)

6.5 Phenology descriptors

★ 6.5.1 Date of vegetative bud break [YYYYMMDD]

When over 50% of terminal buds have enlarged and the bud scales have split exposing the green of the leaves inside

*		6.5.1.1	Days before (-) or after (+) reference standard [d]
*		6.5.1.2 vegetativ	Days of inflorescence bud break before (-) or after (+) ve bud break [d]
*	6.5.2 When 5%	First blo of flowers	oom date [YYYYMMDD] s are opened
*	6.5.3 When 50%	Peak blo 6 of flower	oom date [YYYYMMDD] rs are opened
*		6.5.3.1	Days before (-) or after (+) reference standard [d]
*	6.5.4 When last	Last blo flowers a	om date [YYYYMMDD] re opened
*	6.5.5 When hul	Harvest l separates	date [YYYYMMDD] s easily from the shell
*		6.5.5.1	Days before (-) or after (+) reference standard [d]
	6.5.6	Homoge 0 No 1 Yes	eneity in nut ripening
*	6.5.7 Record wl	Beginni hen trees b	ng of defoliation [YYYYMMDD] begin to defoliate
		6.5.7.1	Days before (-) or after (+) reference standard
*	6.5.8	Defoliati	ion date [YYYYMMDD]

When trees are completely defoliated

6.5.8.1 Days before (-) or after (+) reference standard [d]

6.5.9 Tendency to alternate bearing

Provide an indicative value of the tendency to alternate bearing of the cultivar (e.g. percentage of the production in an off-year in respect to an on-year)

6.6 Notes

Specify here any additional information

EVALUATION

7. Plant descriptors

7.1 Chilling requirements

Number of hours of temperatures below 7°C

- 1 Low (< 600 h)
- 2 Medium (600-1200 h)
- 3 High (> 1200 h)

7.2 Yield

7.2.1 Cropping efficiency [g/cm²]

Nuts yield per unit trunk cross-sectional area. Trunk measurement 20 cm above graft union in grafted tree or 40 cm above ground level in seedling tree

★ 7.2.2 Estimated yield

Rate in relation to age and volume of tree

- 3 Low
- 5 Intermediate
- 7 High

7.3 Kernel

★ 7.3.1 Chemical composition

7.3.1.1	Kernel protein content	[%]
7.3.1.2	Kernel oil content [%]	

- 7.3.1.3 Kernel ash content [%]
- \star

7.3.2 Storage quality

Evaluated 3 months after harvest

★ 7.3.2.1 Kernel rancidity potential [%]
 Polyunsaturated fatty acids content

7.3.2.2 Kernel bitterness

- 3 Weak
- 7 Strong

7.3.2.3 Kernel crispness

- 0 No
- 1 Yes

7.3.2.4 Kernel sweetness 0 No 1 Yes 7.3.2.5 Kernel firmness 0 No 1 Yes

7.4 Pollen

7.4.1 Pollen normal [%]

Incidence of normal grains (normal pollen grains are those \pm equiassic and having an acceptable number and disposition of apertures)

7.4.2 Pollen deformed and/or aborted [%]

Incidence of deformed and/or aborted pollen grains

7.4.3 Ratio of normal/aborted pollen grains

Ratio of the percentages of normal pollen grains over those deformed/aborted

7.4.4 Pollen vitality

Intensity of pollen grain colour after being stained with proline

- 1 Scarcely coloured
- 2 Intensively coloured

7.4.5 Pollen fertility

Intensity of pollen grain colour after being stained with fluorescein

- 1 Scarcely coloured
- 2 Intensively coloured

7.5 Notes

Specify here any additional information

8. Abiotic stress susceptibility

Scored under artificial and/or natural conditions, which should be clearly specified. These are coded on a susceptibility scale from 1 to 9:

- 1 Very low or no visible sign of susceptibility
- 3 Low
- 5 Intermediate
- 7 High
- 9 Very high

8.1 Low temperature

8.1.1 Susceptibility to frost damage in spring

8.2 High temperatures

- 8.2.1 Sunburn susceptibility of hull
- 8.2.2 Sunburn susceptibility of kernel
- 8.2.3 Sunburn susceptibility of trunk

8.3 Salinity

8.4 Mineral deficiency

- 1 Nitrogen
- 2 Phosphorus
- 3 Potassium
- 4 Boron
- 5 Zinc
- 6 Copper
- 99 Other (specify in descriptor 8.8 Notes)

8.5 Mineral toxicity

- 1 Boron
- 2 Zinc
- 3 Chloride
- 4 Copper
- 5 Calcium
- 99 Other (specify in descriptor **8.8 Notes**)

8.6 Waterlogging

8.7 Drought

8.8 Notes

Specify here any additional information

9 Biotic stress susceptibility

In each case, it is important to state the origin of the infestation or infection, i.e. natural, field inoculation, laboratory. Record such information in descriptor **9.9** Notes. These are coded on a susceptibility scale from 1 to 9, viz:

- 1 Very low or no visible sign of susceptibility
- 3 Low
- 5 Intermediate
- 7 High
- 9 Very high

Pistachio bud moth

Thrips

Organisms marked with an asterisk are those found to be of major importance in recent literature sources such as Kaska *et al.* (1995) among others.

Causal organism Taxonomic order or common name 9.1 **Polyphagous insects** 9.1.1 Anapleura lentisci Aphidae 9.1.2 Carpocorus pudicus Heteroptera 9.1.3 Estenoborus persisi Coleoptera 9.1.4 Graphosoma lineatum Heteroptera 9.1.5 Scheidereria pistaciella, S. pistaciicola Lepidoptera 9.1.6 Spilostethus pandurus Pistachio shoot-hole borer 9.1.7 *Sulamicerus stali Cicadellidae 9.1.8 Tinea pistaciae Pistachio shoot-hole borer 9.2 Stem-feeding insects 9.2.1 Capnodis cariosa Coleoptera 9.2.2 Agrillus veridi ceruleus subsp. esfandiarinus Pistachio shoot-hole borer 9.2.3 *Hylesinus vestitus = Acrantus vestitus Pistachio bark beetle, pistachio shoot-hole borer 9.2.4 *Kermania pistaciella Pistachio twig borer 9.3 **Foliage-feeding insects** 9.3.1 *Agonoscena targionii Pistachio psylla 9.3.2 *Anapulvinaria pistaciae Pistachio cushion scale 9.3.3 Ceroplastes rusci Homoptera 9.3.4 *Eulecanium rugulosum Scale Pistachio leafhopper 9.3.5 *Idiocerinus stali 9.3.6 Melanaspis inopinatus Pistachio trunk scale 9.3.7 *Pistaciaspis pistaciae Pistachio scale 9.3.8 Homoptera Saissetia oleae 9.3.9 Slavum wertheimae Gall-forming aphid 9.3.10 *Suturaspis pistaciae Pistachio white scale Pistachio bud moth 9.3.11 *Thaumetopoea solitaria 9.3.12 *Tenuipalpus Pistachio mites 9.3.13 Tetranychus Mites 9.4 **Flower-feeding insects** 9.4.1 Anthascia sp. Coleoptera 9.4.2 Eriophyte pistacia Mites 9.4.3 Frankliniella occidentalis Western flower thrips 9.4.4 Polydrosus davatchi Coleoptera

- **9.4.5** **Telphusa pistaciae*
- **9.4.6** **Thrips iranicus, Thrips pistacia*

9.5 Fruit-feeding insects

9.5.1	*Acrosternum hegeriir	Stink bug
9.5.2	*Amyelois transitella	Naval orangeworm
9.5.3	Apomyelois ceratoniae	Carob moth
9.5.4	Arimania komaroffi Ragonot	Lepidoptera
9.5.5	*Brachynema	Common stink bug
9.5.6	Brevipalpus lewisi	Citrus flat mite
9.5.7	Calocoris norvegicus	Gmelin bug
9.5.8	*Chlorochroa uhleri, C. ligata	Stink bug
9.5.9	Dinarmus pistacia	Hymenoptera
9.5.10	Ephestia elutella	Lepidoptera
9.5.11	*Eurytoma plotnikovi	Pistachio fruit wasp
9.5.12	E. pistachiae	Pistachio wasp
9.5.13	Gonocerus acuteangulatis, Graphosoma semipunctatum	Heteroptera
9.5.14	Leptocoris trivittatus	Boxelder bug
9.5.15	*Leptoglossus clypealis, L. occidentalis	Leaf-footed bug
9.5.16	Liorhyssus hyalinus	Epicarp lesion
9.5.17	Lygaeus hesperus	Hemiptera
9.5.18	Lygaeus panderus	Pistachio red bug
9.5.19	*Megastigmus pistaciae	Pistachio seed chalcid,
	Pistac	chio golden fruit wasp
9.5.20	Neurocolpus californicus, Psallus vaccinola, P. ancorifer	Epicarp lesion
9.5.21	Nezara viridula	Green stink bug
9.5.22	*Plodia interpunctella	Lepidoptera
9.5.23	Phytocoris spp. (Mi	ridae) Epicarp lesion
9.5.24	*Recurvaria pistaciicola	Pistachio fruit moth
9.5.25	Solenostedium bilunatum	
9.5.26	*Thyanta pallidovirens Red-shouldered stink bug	, Pistachio nut worm

9.6 Nematodes

9.6.1	*Heterodera mediterranea, Heterodera marioni	Cyst nematode
9.6.2	Meloidogyne spp.	Root knot nematode
9.6.3	Pratylenchus hamatus, P. neglectus	Root lesion nematode
9.6.4	Xiphinema spp.	Dagger nematode

Rozet virus

9.7 Viruses

(Specify in the descriptor 9.9 Notes)

9.8	Fungi		
	9.8.1	*Alternaria alternata	Alternaria late blight
	9.8.2	Alternaria tenuissima	Alternaria blight
	9.8.3	*Armillaria mellea	Armillaria root rot
	9.8.4	Aspergillus candidus	
	9.8.5	Aspergillus clavatus	

9.8.6	*Aspergillus flavus, A. parasiticus	Seed rot (aflatoxins)
9.8.7	Aspergillus fumigatus	
9.8.8	*Aspergillus ochraceus	Ochratoxins
9.8.9	*Aspergillus niger	
9.8.10	Aspergillus spp.	Aspergillus blights
9.8.11	Asteromella pistaciarum	
9.8.12	*Aureobasidium pullulans	Stigmatomycosis
9.8.13	*Botryosphaeria dothidea	Botryosphaeria panicle and shoot blight
9.8.14	Botryosphaeria obtusa	Stem canker
9.8.15	Botryodiplodia pistaciae	
9.8.16	*Botryotinia fuckeliana (syn. Botryti	<i>is cinerea)</i> Botrytis blossom,
		shoot blight
9.8.17	Cenangium vagabundum	
9.8.18	Camarosporium pistaciae	Camarosporium shoot
9.8.19	Chaetomium spp.	
9.8.20	*Cladosporium herbarum	
9.8.21	Cylindrosporium garbowskii	
9.8.22	C. pistaciae	
9.8.23	Cytospora terebinthi	Gum canker
9.8.24	Cochliobolus spicifer	
9.8.25	*Epicoccum purpurascens	
9.8.26	Eutypa lata	
9.8.27	Fomes rimosus	
9.8.28	Fusarium equiseti	
9.8.29	F. roseum	
9.8.30	F. solani	
9.8.31	*F. oxysporum	
9.8.32	*Fusarium spp.	Root and stem rot
9.8.33	Monilia pistaciae	
9.8.34	Melampsora pistaciae	
9.8.35	Mycosphaerella pistacina	
9.8.36	*Nematospora coryli	Stigmatomycosis
9.8.37	*Paecilomyces variotii	Die-back of young shoots
9.8.38	Pestalotia breviseta.	
9.8.39	*Phytophthora spp.	Gommosis, crown and root rot
9.8.40	Phyllosticta terebinthi	
9.8.41	P. lentisci	
9.8.42	Phymatotrichum omnivorum	Texas root rot
9.8.43	Penicillium camemberti	
9.8.44	P. decumbens	
9.8.45	Penicillium spp.	
9.8.46	*Pestalotiopsis spp.	
9.8.47	Phyllactinia suffulta	
9.8.48	*Phomopsis sp.	Phomopsis shoot blight
		- 0

9.8.49	Pleospora montemartinii	
9.8.50	P. pistaciae	
9.8.51	Pileolaria terebinthi	Rust
9.8.52	Pleurotus ostreatus	
9.8.53	Rhizoctonia bataticola	
9.8.54	Rhizoctonia solani (AG-4)	Nursery seedling blight
9.8.55	Rhizopus sp.	
9.8.56	Rosellina necatrix	
9.8.57	*Schyzophyllum commune S	chyzophyllum wood decay
9.8.58	Sclerotinia sclerotiorum	Sclerotinia shoot blight
9.8.59	Septogloeum pistaciae	
9.8.60	*Septoria pistaciae, S. pistacina (syn. Mycosphar	ella pisticina),
	S. pistaciarum (syn. Mycospherella pistacearum)	Septoria blight
9.8.61	* <i>Septoria</i> spp.	Septoria leaf and fruit spot
9.8.62	Sphaerella pistaciae	
9.8.63	*Stemphyllium botryosum	
9.8.64	<i>Tetracoccosporium</i> sp.	
9.8.65	Trichoderma harzianum	
9.8.66	*Trichothecium roseum	
9.8.67	Tzavella roumbos	Panicle blight
9.8.68	Uromyces terebinthi (syn. Pileolaria terebinthi)	Rust
9.8.69	*Verticillium albo-atrum, V. dahliae	Verticillium wilt

9.9 Notes

Specify here any additional information

10. Molecular markers

Describe any specific discriminating or useful trait for this accession. Report probe-enzyme combination analyzed. Below are listed some of the basic methods most commonly used

10.1 Restriction fragment length polymorphism (RFLP)

Report probe/enzyme combination (approach can be used for nuclear, chloroplast or mitochondrial genomes)

10.2 Amplified fragment length polymorphism (AFLP)

Report primer pair combinations and accurate molecular size of products (used for nuclear genomes)

10.3 DNA amplification fingerprinting (DAF); random amplified polymorphic DNA (RAPD); AP-PCR

Accurately report experimental conditions and molecular size of products (used for nuclear genomes)

10.4 Sequence-tagged microsatellites (STMS)

Report primer sequences, and accurate product sizes (can be used for nuclear or chloroplast genomes)

10.5 PCR-sequencing

Report PCR primer sequences, and derived nucleotide sequence (can be used for single copy nuclear, chloroplast or mitochondrial genomes)

10.6 Other molecular markers

11. Cytological characters

11.1 Chromosome number

11.2 Ploidy level

(2x, 3x, 4x, etc.)

11.3 Meiosis chromosome associations

Average of 50 microspore mother cells, observed during metaphase 1

11.4 Other cytological characters

12. Identified genes

Describe any known specific mutant present in the accession

REFERENCES

- FAO. 1990. Guidelines for Soil Profile Description, 3rd edition (revised). Food and Agriculture Organization of the United Nations, International Soil Reference Information Centre, Land and Water Development Division. FAO, Rome.
- Ferguson, L. (ed.) 1995. Pistachio production. Centre for Fruit and Nut Crop Research and Information. University of California Davies, California, USA, 126p.
- Kaska, N., A.B. Küden, L. Ferguson, and T. Michailides (eds.). 1995. Pistachio Nut I. Acta Horticulturae. 425p.
- Kornerup, A. and J.H. Wanscher. 1984. Methuen Handbook of Colour. Third edition. Methuen, London.
- Munsell Color. 1977. Munsell Color Charts for Plant Tissues, 2nd edition, revised. Munsell Color, Macbeth Division of Kollmorgen Corporation, 2441 North Calvert Street, Baltimore, Maryland 21218, USA.
- Padulosi, S., T. Caruso and E. Barone (eds.) 1996. Taxonomy, distribution, conservation and uses of *Pistacia* genetic resources. Report of a workshop, 29-30 June 1995. Palermo, Italy. International Plant Genetic Resources Institute, Rome, Italy, 69p.
- Rana, R.S., R.L. Sapra, R.C. Agrawal and Rajeev Gambhir. 1991. Plant Genetic Resources. Documentation and Information Management. National Bureau of Plant Genetic Resources (Indian Council of Agricultural Research). New Delhi, India.
- Royal Horticultural Society. 1966, c. 1986. R.H.S. Colour Chart (ed. 1, 2). Royal Horticultural Society, London.
- Sheibani, A. 1990. Characteristics of pistachio selected rootstocks. Nahal-O-Bazr, Research Journal. Seed and Plant Improvement Inst., Karaj. Iran. Vol 6: (1,2) 49-59.
- Sheibani, A. 1994. Pistachio production in Iran. Pistachio Research Institute, Rafsanjan, Iran.
- van Hintum, Th.J.L. 1993. A computer compatible system for scoring heterogeneous populations. Genetic Resources and Crop Evolution 40:133-136.
- Venturella, G. 1991. A check-list of Sicilian fungi, Bocconea 2:5-221.
- Zohary, M. 1952. A monographic study of the genus *Pistacia*. Palestine J. Bot., Jerusalem series 5(4):187-228.

CONTRIBUTORS

Authors

Dr Ettore Barone Istituto di Coltivazioni Arboree Faculty of Agriculture University of Palermo Viale delle Scienze 90128, Palermo ITALY Tel. (+39) 91-423398 Fax. (+39) 91-6521098 email ebarone@mbox.unipa.it

Dr Stefano Padulosi IPGRI Via delle Sette Chiese, 142 00145, Rome ITALY Tel. (+39) 6-51892243 Fax (+39) 6-5750309 email s.padulosi@cgnet.com

Ir Paul Van Mele University of Ghent Faculty of Agricultural and Applied Biological Sciences Lab. Tropical and Sub-Tropical Agriculture and Ethnobotany Dept. of Plant Production Coupure Links, 653B-9000 Ghent BELGIUM Tel. (+32) 9-2646089 Fax. (+32) 9-2646241 email paul.vanmele@rug.ac.be

Reviewers

Prof. Bekir Erol Ak Harran University Faculty of Agriculture Dept. of Horticulture Sanliurfa TURKEY Tel. (+90) 414- 2470383 Fax. (+90) 414- 2470385

Dr Ignacio Batlle IRTA Centre de Mas Bové Departament d'Arboricultura Mediterrània Apartat 415, E-43280 Reus SPAIN Tel. (+34) 77-343252 Fax. (+34) 77-344055 email ignasi@masbove.irta.es

Prof Tiziano Caruso Istituto di Coltivazioni Arboree Faculty of Agriculture University of Naples Via Università 100 80055, Portici (NA) ITALY Tel. (+39) 81-7755141 Fax. (+39) 81-7755114

Prof Francesco Giulio Crescimanno Istituto di Coltivazioni Arboree Faculty of Agriculture University of Palermo Viale delle Scienze 90128, Palermo ITALY Tel. (+39) 91-423398 Fax. (+39) 91-6521098 Dr Laura De Palma Istituto di Coltivazioni Arboree Faculty of Agriculture University of Bari Via Amendola 165 70126, Bari ITALY Tel. (+39) 80-5442982 Fax. (+39) 80-5442813

Prof Luigi Di Marco Istituto di Coltivazioni Arboree Università degli Studi di Palermo Facoltà de Scienze Agrarie Viale delle Scienze 11 90128, Palermo ITALY Tel. (+39) 91-423398 Fax. (+39) 91-6521098

Prof Ahmed El-Oqlah Dept. of Biological Sciences Faculty of Science University of Yarmouk Irbid JORDAN Tel. (+962) 2-271100 ext. 2831 Fax. (+962) 2-274725

Prof Louise Ferguson Dept. of Pomology University of California Kearney Agricultural Center 9420 South Riverbend Dr. Parlier, CA 93648 USA Tel. (+1) 209-8912500 Fax. (+1) 209-8912593 email louise@uckac.edu Prof Ruhinaz Gulcan Ege University Faculty of Agriculture Dept. of Horticulture 35100 Bornova- Izmir TURKEY Tel. (+90) 232-3880110 Fax. (+90) 232-3881864

Prof Ahamad Hadj-Hassan Aleppo University Faculty of Agriculture Dept. of Horticulture 021 Aleppo SYRIA Fax. (+963) 21- 213490

Prof Abdallah Jaradat IPGRI c/o ICARDA PO Box 5466 Aleppo, SYRIA Tel. (+963) 21-247485 Fax. (+963) 21-225105/213490 email a.jaradat@cgnet.com

Prof Nurettin Kaska Dept. of Horticulture Faculty of Agriculture University of Çukurova 01330, Adana TURKEY Tel. (+90) 322-3386748 Fax. (+90) 322-3386388

Dr Abdellamid Khaldi Institut National de Recherches Forestières BP 2, 2080 Ariana TUNISIA Tel. (+216) 1-230420 Fax. (+216) 1-750557 Prof Mohamed Laghezali Programme d' Arboriculture Fruitiere CRSMA, 3 Esplanade du Dr Giguet BP 578, Meknès MOROCCO Tel. (+212) 5-520300 Fax. (+212) 5-515953/512040

Prof Pietro Mazzola Dipartimento di Scienze Botaniche Faculty of Agriculture University of Palermo Via Archirafi 38 90123, Palermo ITALY Tel. (+39) 91-6161493 Fax. (+39) 91-6176089

Prof Themis J. Michailides Dept. of Pomology University of California Kearney Agricultural Center 9420 South Riverbend Drive Parlier, CA 93648 USA Tel. (+1) 209-8912500 Fax. (+1) 209-8912593

Dr Mohamed Mlika INRAT Institut National de la Recherche Agronomique Laboratoire d'Arboriculture Fruitière Avenue de l' Indépendance Ariana TUNISIA Tel. (+216) 1- 230024 Fax. (+216) 1- 231693 Prof Francesco Monastra Istituto Sperimentale per la Frutticoltura Via Fioranello 52 00040 Ciampino Aeroporto Roma ITALY Tel. (+39) 6- 79340251 Fax. (+39) 6- 79340158

Prof Vittorino Novello Istituto di Coltivazioni Arboree Faculty of Agriculture University of Bari Via Amendola 165 70126, Bari ITALY Tel. (+39) 80- 5442982 Fax. (+39) 80- 5442813

Prof Francesco M. Raimondo Istituto di Botanica Faculty of Agriculture University of Palermo Viale delle Scienze 90128, Palermo ITALY Tel. (+39) 91- 6161493 Fax. (+39) 91- 617689

Dr Miguel A. Romero IRTA Centre de Mas Bové Departament d'Arboricultura Mediterrània Apartat 415 E-43280 Reus SPAIN Tel. (+34) 77- 343252 Fax. (+34) 77- 344055 email lannoye@masbove.irta.es Dr Dimos Rouskas NARF National Agricultural Research Foundation Vardates Station N. Krikello G-35100 Lamias GREECE Tel. (+30) 2- 3181246/ 81104 Fax. (+30) 2- 3181822

Dr Ahmed Sheibani Pistachio Research Institute PO Box 77375-435 Rafsanjan IRAN Tel. (+98) 3431- 2062/3026 Fax. (+98) 3431- 4611

Prof Patrick Van Damme University of Ghent, Faculty of Agricultural and Applied Biological Sciences Lab. Tropical and Sub-Tropical Agriculture and Ethnobotany Dept. of Plant Production Coupure Links 653B-9000 Ghent BELGIUM Tel. (+32) 9 - 2646089 Fax. (+32) 9 - 264 6241 email patrick.vandamme@rug.ac.be

Dr Francisco J. Vargas IRTA Centre de Mas Bové Apartado 415 43280 Reus SPAIN Tel. (+34) 77- 343252 Fax. (+34) 77- 344055 email vargas@masbove.irta.es Prof Giuseppe Venturella Dept. Botanic Sciences Faculty of Agriculture University of Palermo Via Archirafi 38 90123, Palermo ITALY Tel. (+39) 91- 6161493 Fax. (+39) 91- 6176089

Dr Georgios Zakintinos NARF National Agricultural Research Foundation Vardates Station N. Krikello G-35100 Lamias GREECE Tel. (+30) 2- 3181246/ 81104 Fax. (+30) 2- 3181822

Prof Daniel Zohary Dept. of Evolution, Systematic and Ecology Hebrew University 91904 - Jerusalem ISRAEL Tel. (+972) 2- 792927/ 6585390 Fax. (+972) 2- 792613

ACKNOWLEDGEMENTS

IPGRI wishes to place on record their sincere thanks to the numerous pistachio workers around the world who have contributed directly or indirectly to the development of Descriptors for Pistachio. IPGRI is particularly grateful to Prof Luigi di Marco, University of Palermo and Prof Tiziano Caruso of the University of Naples for having kindly provided support and scientific advice during the development phase of the descriptors. National pistachio descriptors lists of Iran, Turkey and Spain (Istituto di Ricerca i Tecnologia Agroalimentare) have been used as valuable references for the preparation of this document.

Ms Adriana Alercia supervised and coordinated the production of the text up to the prepublication stage and provided technical expertise. Ms Linda Sears edited the text, and Ms Patrizia Tazza designed the cover and prepared the layout. Dr S. Padulosi designed the illustrations. Mr Paul Stapleton managed the production of the publication. Ir Tom Hazekamp provided scientific direction and supervised the overall process.

The following IPGRI staff provided substantial technical advice: Drs. Marlene Diekmann and Toby Hodgkin.

ANNEX I. MULTICROP PASSPORT DESCRIPTORS

This list of multicrop passport descriptors has been developed jointly by IPGRI and FAO to provide consistent coding schemes for common passport descriptors across crops. These descriptors aim to be compatible with future IPGRI crop descriptor lists and with the descriptors to be used for the FAO World Information and Early Warning System (WIEWS) on plant genetic resources.

The list should NOT be regarded as a minimum descriptor list, since many additional passport descriptors are essential for the description of crops and need to be recorded. This document lists an initial set of common passport descriptors at the multicrop level. At a later stage the list could be expanded with additional multicrop descriptors. For example, descriptors dealing with the use of germplasm are currently not included, but their suitability for inclusion at the multicrop level will be investigated. Future expansion could even result in the development of more specialized lists of common descriptors at the crop group level.

Printed here is the latest version of the list (1997) which contains two sections. The latter one (FAO WIEWS DESCRIPTORS) lists a number of optional descriptors used in the FAO WIEWS. The list provides descriptions of content and coding schemes, but also provides *suggested* fieldnames (in parentheses) that can assist in the computerized exchange of this type of data.

MULTICROP PASSPORT DESCRIPTORS

1. Institute code

Code of the institute where the accession is maintained. The codes consist of the 3-letter ISO 3166 country code of the country where the institute is located plus number or an acronym as specified in the Institute database that will be made available by FAO. Preliminary codes (i.e. codes not yet incorporated in the FAO Institute database) start with an asterisk followed by a 3letter ISO 3166 country code and an acronym.

2. Accession number

This number serves as a unique identifier for accessions and is assigned when an accession is entered into the collection. Once assigned this number should never be reassigned to another accession in the collection. Even if an accession is lost, its assigned number should never be reused. Letters should be used before the number to identify the genebank or national system (e.g. IDG indicates an accession that comes from the genebank at Bari, Italy; CGN indicates an accession from the genebank at Wageningen, The Netherlands; PI indicates an accession within the USA system).

Collecting number 3.

Original number assigned by the collector(s) of the sample, normally composed of the name or initials of the collector(s) followed by a number. This item is essential for identify-ing duplicates held in different collections. It should be unique and always accompany subsamples wherever they are sent.

4 Genus

Genus name for taxon. Initial uppercase letter required.

Species 5.

Specific epithet portion of the scientific name in lowercase letters plus authority¹. Following abbreviation is allowed: "sp."

Subtaxa 6.

Subtaxa can be used to store any additional taxonomic identifier plus authority¹. Following abbreviations are allowed: "ssp." (for subspecies); "var." (for variety); "convar." (for convariety); "f." (for form).

7. Accession name

(ACCNAME) Either a registered or other formal designation given to the accession. First letter uppercase. Multiple names separated with semicolon.

Country of origin 8.

Name of the country in which the sample was originally collected or derived. Use the ISO 3166 extended codes, (i.e. current and old 3 letter ISO 3166 country codes)

9. Location of collecting site

Location information below the country level that describes where the accession was collected starting with the most detailed information. Might include the distance in kilometers and direction from the nearest town, village or map grid reference point, (e.g. CURITIBA 7S, PARANA means 7 km south of Curitiba in the state of Parana)

¹ Authority is only provided at the most detailed taxonomic level

(INSTCODE)

(ACCENUMB)

(COLLNUMB)

(SPECIES)

(ORIGCTY)

(COLLSITE)

(GENUS)

(SUBTAXA)

10. Latitude of collecting site

(LATITUDE) Degrees and minutes followed by N (North) or S (South) (e.g. 1030S). Missing data (minutes) should be indicated with hyphen (e.g. 10—S).

11. Longitude of collecting site (LONGITUDE) Degrees and minutes followed by E (East) or W (West) (e.g. 07625W). Missing data (minutes) should be indicated with hyphen (e.g. 076–W).

12. Elevation of collecting site [m asl]

Traditional cultivar/Landrace

(ELEVATION) Elevation of collecting site expressed in meters above sea level. Negative values allowed.

13. Collecting date of original sample [YYYYMMDD]

(COLLDATE) Collecting date of the original sample where YYYY is the year, MM is the month and DD is the day.

14.	Status	of	sample
1	3 3 7 1 1		-

Wild 1 2 Weedy

3

0 Unknown (SAMPSTAT)

99 Other (Elaborate in REMARKS field)

4 Breeder's line Advanced cultivar 5

15. Collecting source

(COLLSRC) The coding scheme proposed can be used at 2 different levels of detail: Either by using the global codes such as 1, 2, 3, 4 or by using the more detailed coding such as 1.1, 1.2, 1.3 etc.

Wild habitat	2	Farm	3	Market	4	Institute/ Research
1.1 Forest/woodland		2.1 Field		3.1 Town		organization
1.2 Shrubland		2.2 Orchard		3.2 Village		0
1.3 Grassland		2.3 Garden		3.3 Urban	0	Unknown
1.4 Desert/tundra		2.4 Fallow		3.4 Other	9	90ther
		2.5 Pasture		exchange	. (Elaborate in
		2.6 Store		system	À	EMARKS field)

16. Donor institute code

Code for the donor institute. The codes consist of the 3-letter ISO 3166 country code of the country where the institute is located plus number or an acronym as specified in the Institute database that will be made available by FAO. Preliminary codes (i.e. codes not yet incorporated in the FAO Institute database) start with an asterisk followed by a 3-letter ISO 3166 country code and an acronym.

17. Donor number

Number assigned to an accession by the donor. Letters should be used before the number to identify the genebank or national system (e.g. IDG indicates an accession that comes from the genebank at Bari, Italy; CGN indicates an accession from the genebank at Wageningen, The Netherlands; PI indicates an accession within the USA system).

18. Other number(s) associated with the accession

Any other identification number known to exist in other collections for this accession. Letters should be used before the number to identify the genebank or national system (e.g. IDG indicates an accession that comes from the genebank at Bari, Italy; CGN an accession from the genebank at Wageningen, The Netherlands; PI indicates an accession within the USA system). Multiple numbers can be added and should be separated with a semicolon.

(DONORNUMB)

(OTHERNUMB)

(DONORCODE)

19. Remarks

(REMARKS)

The remarks field is used to add notes or to elaborate on descriptors with value "99" (=Other). Prefix remarks with the field name they refer to and a colon (e.g. COLLSRC: roadside). Separate remarks referring to different fields are separated by semicolons.

FAO WIEWS DESCRIPTORS

1. Location of safety duplicates

Code of the institute where a safety duplicate of the accession is maintained. The codes consist of 3-letter ISO 3166 country code of the country where the institute is located plus number or an acronym as specified in the Institute database that will be made available by FAO. Preliminary codes (i.e. codes not yet incorporated in the FAO Institute database) start with an asterisk followed by a 3-letter ISO 3166 country code and an acronym. Multiple numbers can be added and should be separated with a semicolon.

2. Availability of passport data (i.e. in addition to what has been pro	ovided)		(PASSAVAIL)
0 Not available	1	Available	
3. Availability of characterizatio	n data		(CHARAVAIL)
0 Not available	1	Available	. , ,
4. Availability of evaluation data	1		(EVALAVAIL)
0 Not available	1	Available	
5. Acquisition type of the acces	sion		(ACQTYPE)
1 Collected/bred originally by th	e institut	e	
2 Collected/bred originally by jo	int missio	on/institution	
3 Received as a secondary reposi	tory		
6. Type of storage			(STORTYPE)
Maintenance type of germplasm. If g	germplas	m is maintained unde	er different types of stor-
age, multiple choices are allowed, se	parated b	y a semicolon (e.g. 2;	3). (Refer to FAO/IPGRI
Genebank Standards 1994 for details	s on stora	ge type)	
1 Short-term	5	Field genebank col	lection
2 Medium-term	6	Cryopreserved	
3 Long-term	99	Other (elaborate in	REMARKS field)
4 In vitro collection			

Please forward your feedback on the use of this list to: Tom Hazekamp, Germplasm Documentation Officer International Plant Genetic Resources Institute

Via delle Sette Chiese 142 00145 Rome, Italy Email: T.HAZEKAMP@CGNET.COM Fax: (+39) 065750309

(DUPLSITE)

COLLECTING FORM for pistachio (Pistacia vera L.)

ACCESSION No. (1.1):

COLLECTOR NAME(S)/INSTITUTE(S) (2.1):

ACCESSION IDENTIFICATION	
COLLECTING No. (2.3):	 []
COLLECTING DATE (YYYY/MM/DD) (2.4):	
GENUS (1.8.1): SPECIES	(1.8.3):
SEX (1.11): 1 Male 2 Female	
LOCAL/VERNACULAR NAME (2.16):	
ETHNIC GROUP (2.17):	
LOCAL LANGUAGE:	
ORCHARD MANAGEMENT	
ACCESSION NUMBER (3.1)	
TYPE OF MAINTENANCE (3.10) 1. Vegetative in the field 2. Vegetative in tissue cultu 99. Other (specify):	re 3. Pollen 4. Seed
CHARACTERIZATION	
LEAF DESCRIPTORS (6.2) Terminal leaflet length [cm] (6.2.4): Terminal leaflet width [cm] (6.2.5):	
INFLORESCENCE AND FRUITING HABIT (6.3) Female reference standard (6.3.1): Male reference standard (6.3.2): Flowering precocity (6.3.3):	
NUT AND KERNEL (6.4) Nut length [mm] (6.4.6): Nut width [mm] (6.4.7): Nut thickness (6.4.8): Nut shape (6.4.9): Shell apex (6.4.10):	Suture opening (6.4.20): Blank production (6.4.22): Kernel colour (6.4.32): Festa colour (6.4.33): Estimated yield (7.2.2)
SAMPLE	
TYPE OF SAMPLE (2.13): 1. Vegetative 2. Seed 3. Pollen 4	ł. Tissue culture

STATUS OF SA 0. Unknown 4.Breeders line	MPLE (2.14) 1.Wild 5.Advanced c	ultivar	2.Weed 99.Othe	y er (speci	fy):	3.Traditional cultivar/Landrace					
USES OF THE 1.Nut production 5.Medicinal	FRUIT (2.22) 1 2.Clonal roots 6.Forage	tock	3.Seedl 7.Wood	ing root I/timber	stock	4. Pollinator 99.Other (specify):					
PREVAILING STRESSES (2.27): Mention the types of major stresses i.e. abiotic (drought), biotic (insects, fungi, etc.)											
COLLECTING SITE LOCATION											
 COUNTRY. (2.5): _ _ _ _ _ _ _ REGION: _											
PROVINCE/STA	OVINCE/STATE (2.6): DEPARTMENT/COUNTY (2.7):										
_OCATION (2.8) km: direction: from:											
LATITUDE (2.9): _ _ _ _ _ _ LONGITUDE (2.10): _ _ _ _ _ _ _ _ _											
ACCESSION A		IG SITE	ENVIRO	NNEME	NT						
COLLECTING S 0.Unknown 4.Institute/Rese	OURCE (2.12): 1.Wild habitat arch organizatio	n	2.Farm 99.Othe	r (speci	fy):	3.Marke	et				
HIGHER LEVEL LANDFORM (5.1.2) 1.Plain 2.Basin 5.Upland 6.Hill		5.1.2)	3.Valley 4.Plateau 7.Mountain								
SLOPE [°](5.1.4): SOIL FERTILITY (5.1.20): (code: 3=Low ; 5=Moderate ; 7=High)											
SOIL TEXTURE (State class (e.g	CLASSES (5.1 Clay, Loam, S	.17) ilt)									
SOIL TAXONON State class (e.g.	/IC CLASSIFIC Alfisols, Spodo	ATION (5 sols, Ver	5.1.18) tisols)								
WATER AVAILABILITY (5.1.19) 1.Rainfed 2.Irrigated 5.Sea coast 99.Other (specify):		cify):	3.Flooded			4.River banks					
RAINFALL (5.1. JAN F Seasonal: mean (mm)	21.2) FEB MAR / 	Annual I APR I I I	mean: _ MAY I	 JUN 	mm JUL II	AUG II	SEP 	ОСТ II	NOV 	DEC	
TEMPERATURE JAN F Seasonal: mean (°C)	E (5.1.21.1) FEB MAR /	Annual m APR I I I	iean: MAY I	°(JUN 	C JUL I_I	AUG	SEP 	OCT 	NOV 	DEC	

53