



Trigonella foenum-graecum Seed – Identification

Thin-Layer Chromatography – Amino Acids Profile



A



B


Typical HPTLC Chromatograms

These chromatograms are supplied for information only


Track assignment: 1-3) *Trigonella foenum-graecum* Seed, commercial samples; 4) USP 4-Hydroxyisoleucine RS (0.5 mg/mL); 5-13) finished products (capsules and tablets); 14-16) liquid extracts

Sample solutions:	according to the monograph
Standard solutions:	in methanol
Plate:	HPTLC, Si 60 F ₂₅₄
Saturation time:	20 minutes
Application volume:	2 µL, as 8-mm bands
Relative Humidity:	about 33%
Temperature:	25°
Developing solvent system:	<i>n</i> -Butanol, acetic acid, and water (7:2:1)
Developing distance:	6 cm
Derivatization reagent:	ninhydrin reagent – 0.3 g of ninhydrin, 95 mL of isopropanol, and 5 mL of glacial acetic acid
Detection:	derivatize, heat at 100-105° for 2 min, and examine under (A) UV light at 366 nm and (B) visible light.

Thin-Layer Chromatography – Steroidal Saponins Profile



A



B

Typical HPTLC Chromatograms


These chromatograms are supplied for information only

Track assignment: 1-3) *Trigonella foenum-graecum* Seed, commercial samples;
 4) fructose, protogracillin, protodioscin and dioscin (two bands), 0.5 mg each/mL (with increasing R_f);
 5-13) finished products (capsules and tablets); 14-16) liquid extracts.

Sample solutions:	according to the monograph
Standard solutions:	in methanol
Plate:	HPTLC, Si 60 F ₂₅₄
Saturation time:	20 minutes
Application volume:	2 μ L, as 8-mm bands
Relative Humidity:	about 33%
Temperature:	25°
Developing solvent system:	chloroform, methanol and water (18:8:1)
Developing distance:	6 cm
Derivatization reagent:	anisaldehyde reagent – 85 mL of ice-cooled methanol mixed with 10 mL of glacial acetic acid, 5 mL of sulfuric acid, and 0.5 mL of <i>p</i> -anisaldehyde

Detection: derivatize, heat at 100° for 2-3 min, and examine under (A) UV light at 366 nm and (B) visible light

Thin-Layer Chromatography – Presence of Trigonelline



Typical HPTLC Chromatograms

These chromatograms are supplied for information only

Track assignment: 1-3) *Trigonella foenum-graecum* Seed, commercial samples; 4) USP Trigonelline Hydrochloride RS (1.5 mg/mL); 5-13) finished products (capsules and tablets); 14-16) liquid extracts.

Sample solutions: according to the monograph

Standard solutions: in methanol

Plate: HPTLC, Si 60 F₂₅₄

Saturation time: 20 minutes

Application volume: 4 µL, as 8-mm bands

Relative Humidity: about 33%

Temperature: 25°

Developing solvent system: isopropyl alcohol, methanol, and water (4:1:4)

Developing distance: 6 cm

Detection: dry, and examine under UV light at 254 nm